

References

- Abbott, M. B., Bathurst, J. C., Cunge, J. A., O'Connell, P. E. and Rasmussen, J. (1986). An introduction to the European Hydrologic System – Système Hydrologique Européen, “SHE”, 1, History and philosophy of a physically-based, distributed modelling system. *J. Hydrology*, 87: 45–9.
- Abbott, M. B. and Refsgaard, J. C. (1996). *Distributed Hydrological Modelling*. Kluwer, Dordrecht, The Netherlands, 321pp.
- Abrahams, A. D., Parsons, A. J. and Luk, S. H. (1989). Distribution of depth of overland flow on desert hillslopes and its implications for modelling soil erosion. *J. Hydrology*, 106: 177–84.
- Ahmed, S. and de Marsily, G. (1987). Comparison of geostatistical methods for estimating transmissivity using data on transmissivity and specific capacity. *Water Resour. Res.*, 23: 1717–37.
- Air Weather Service (1979). *The Use of the Skew T, Log P Diagram in Analysis and Forecasting*. Air Weather Service Manual, Illinois.
- Allard, D. (1993). On the connectivity of two random set models: the truncated Gaussian and the Boolean. In: Soares, A. (Ed.), *Geostatistics Tróia '92. Quantitative Geology and Geostatistics*. Kluwer, Dordrecht, The Netherlands, pp.467–78.
- Allard, D. (1994). Simulating a geological lithofacies with respect to connectivity information using the truncated Gaussian model. In: Armstrong, M. and Down, P. A. (Eds), *Geostatistical Simulations*. Proc. Geostatistical Simulation Workshop, Fontainebleau, France, 27–28 May 1993. *Quantitative Geology and Geostatistics*. Kluwer, Dordrecht, The Netherlands, pp.197–211.
- Altse, E., Bolognani, O., Mancini, M. and Troch, P. A. (1996). Retrieving soil moisture over bare soil from ERS-1 synthetic aperture radar data: Sensitivity analysis based on a theoretical surface scattering model and field data. *Water Resour. Res.*, 89(3): 653–61.
- Ambrose, B., Beven, K. and Freer, J. (1996). Toward a generalization of the TOPMODEL concepts: topographic indices of hydrological similarity. *Water Resour. Res.*, 32(7): 2135–45.
- Anderson, E. A. (1973). National Weather Service River Forecast System–Snow Accumulation and Ablation Model, NOAA Technical Memorandum NWS HYDRO-17, U.S. Dept. of Commerce.
- Anderson, E. A. (1976). A Point Energy and Mass Balance Model of a Snow Cover, NOAA Technical report NWS 19, U.S. Dept. of Commerce.

- Anderson, M. C., Norman, J. M., Diak, G. R., Kustas, W. P. and Mecikalski, J. R. (1997). A two source time-integrated model for estimating surface fluxes from thermal infrared satellite observations. *Remote Sens. Environ.*, 60: 195–216.
- Anderson, M. G. and Burt, T. P. (1978a). Experimental investigations concerning the topographic control of soil water movement on hillslopes. *Z. Geomorph. N. F.*, Suppl. Bd. 29: 52–63.
- Anderson, M. G. and Burt, T. P. (1978b). The role of topography in controlling throughflow generation. *Earth Surface Processes*, 3: 331–44.
- Anderson, M. G. and Burt, T. P. (1978c). Toward more detailed field monitoring of variable source areas. *Water Resour. Res.*, 14(6): 1123–31.
- Anderson, M. G. and Kneale, P. E. (1980). Topography and hillslope soil water relationships in a catchment of low relief. *J. Hydrology*, 47: 115–28.
- Anderson, M. G. and Kneale, P. E. (1982). The influence of low-angled topography on hillslope soil-water convergence and stream discharge. *J. Hydrology*, 57: 65–80.
- Anderson, M. G. and Rogers, C. C. M. (1987). Catchment scale distributed hydrological models – A discussion of research directions. *Progress in Physical Geography*, 11: 28–51.
- Anderson, M. P. (1997). Characterization of geological heterogeneity. In: Dagan, G. and Neuman, S. P. (Eds), *Subsurface Flow and Transport*. Cambridge University Press, pp. 23–43.
- Anderson, M. P. and Woessner, W. W. (1992). The role of postaudit in model validation. *Adv. Water Resour.*, 15: 167–73.
- Armstrong, B. and Williams, K. (1986). *The Avalanche Book*, Fulcrum Inc, Golden, Colorado, 256pp.
- Armstrong, M. (1998). *Basic Linear Geostatistics*. Springer, Berlin.
- Attema, E. (1991). The active microwave instrument on-board the ERS-1 satellite. *Proc. IEEE*, 79: 791–9.
- Austin, P. M. and Houze, R. A. (1972). Analysis of the structure of precipitation patterns in New England. *J. Appl. Meteor.*, 11: 926–35.
- Avissar, R. (1992). Conceptual aspects of a statistical-dynamical approach to represent landscape subgrid-scale heterogeneities in atmospheric models. *J. Geophys. Res.*, 97: 2729–42.
- Avissar, R. (1995). Scaling of land–atmosphere interactions: An atmospheric modelling perspective. *Hydrological Processes*, 9: 679–95.
- Avissar, R. (1998). Which type of soil-vegetation-atmosphere transfer scheme is needed for general circulation models: a proposal for a higher-order scheme. *J. Hydrology*, 212–13: 136–54.
- Avissar, R. and Liu, Y.-Q. (1996). Three dimensional numerical study of shallow convective clouds and precipitation induced by land surface forcing. *J. Geophys. Res.*, 101(D3): 7499–518.
- Bair, E. S. (1994). Model (in)validation – a view from the courtroom. *Ground Water*, 32(4): 530–1.
- Baldocchi, D. D. (1993). Scaling water vapor and carbon dioxide exchange from leaves to a canopy: rules and tools. In: Ehleringer, J. R. and Field, C. B. (Eds), *Scaling Physiological Processes Leaf to Globe*. Academic Press, New York, pp.77–114.
- Baldocchi, D., Hicks, B. B. and Meyers, T. P. (1988). Measuring biosphere–atmosphere exchanges of biologically related gases with micrometeorological methods. *Ecology*, 69(5): 1331–40.
- Band, L. E. and Moore, I. D. (1995). Scale: Landscape attributes and Geographical Information Systems. *Hydrological Processes*, 9: 401–22.
- Band, L. E., Patterson P., Nemani, R. and Running, S. (1993). Forest ecosystem processes at the watershed scale: incorporating hillslope hydrology. *Agric. For. Meteorol.*, 63: 93–126.
- Bárdossy, A. and Lehmann, W. (1998). Spatial distribution of soil moisture in a small catchment. Part 1: geostatistical analysis. *J. Hydrology*, 206: 1–15.

- Barling, R. D., Moore, I. D. and Grayson, R. B. (1994). A quasi-dynamic wetness index for characterising the spatial distribution of zones of surface saturation and soil water content. *Water Resour. Res.*, 30: 1029–44.
- Barros, A. P. and Lettenmaier, D. P. (1993). Dynamic modelling of the spatial distribution of precipitation in remote mountainous areas. *Monthly Weather Review*, 121(April): 1195–214.
- Barros, A. P. and Lettenmaier, D. P. (1994). Dynamic modelling of orographically induced precipitation. *Rev. Geophys.*, 32(3): 265–84.
- Bastiaanssen, W. G. M., Feddes, R. A. and Holtslag, A. A. M. (1998). A remote sensing surface energy balance algorithm for land (SEBAL) Part 1: Formulation. *J. Hydrology*, 212–213(1–4): 198–212.
- Bathurst, J. C. and O'Connell, P. E. (1992). Future of distributed modelling: The Système Hydrologique Européen. *Hydrological Processes*, 6: 265–77.
- Bathurst, J. C., Wicks, J. M. and O'Connell, P. E. (1995). The SHE/SHESED basin scale water flow and sediment transport modelling system. In: Singh, V. P. (Ed.), *Computer Models of Watershed Hydrology*. Water Resources Publications, Highlands Ranch, Colorado, pp. 563–94.
- Beck, M. B. (1987). Water quality modeling: a review of the analysis of uncertainty. *Water Resour. Res.*, 23: 1393–442.
- Becker, A., Blöschl, G. and Hall, A. (1999). Land surface heterogeneity and scaling in hydrology. *J. Hydrology* (special issue), 217(3–4).
- Bergström, S. (1991). Principles and confidence in hydrological modelling. *Nordic Hydrology*, 22: 123–36.
- Betson, R. P. (1964). What is watershed runoff? *J. Geophys. Res.*, 69(8): 1541–52.
- Betson, R. P. and Ardis, C. V. (1978). Implications for modelling surface-water hydrology. Chapter 8 in: Kirkby, M. J. (Ed.), *Hillslope Hydrology*. Wiley, Chichester, pp.295–323.
- Beven, K. J. (1986). Runoff production and flood frequency in catchments of order n : an alternative approach. In: Gupta, V. K., Rodriguez-Iturbe, I. and Wood, E. F. (Eds), *Scale Problems in Hydrology*. Reidel, Dordrecht, pp.107–31.
- Beven, K. J. (1987). Towards a new paradigm in hydrology. In: *Water for the future: Hydrology in perspective*. IAHS Publ. No. 164: 393–403.
- Beven, K. (1989). Changing ideas in hydrology – the case of physically-based models. *J. Hydrology*, 105: 157–72.
- Beven, K. (1995). Linking parameters across scales: Subgrid parameterizations and scale dependent hydrological models. *Hydrological Processes*, 9: 507–25.
- Beven, K. J. (1996). A discussion of distributed hydrological modelling. In: Abbott, M. B. and Refsgaard, J. C. (Eds), *Distributed Hydrological Modelling*, Kluwer, Dordrecht, The Netherlands, pp. 255–78.
- Beven, K. (1997). TOPMODEL: A critique. *Hydrological Processes*, 11: 1069–85.
- Beven, K. J. and Binley, A. M. (1992). The future of distributed models: model calibration and uncertainty prediction. *Hydrological Processes*, 6: 279–98.
- Beven, K. J. and Fisher, J. (1996). Remote sensing and scaling in hydrology. In: Stewart, J. B., Engman, E. T., Feddes, R. A. and Kerr, Y. (Eds), *Scaling up in Hydrology using Remote Sensing*. Wiley, London, pp.1–18.
- Beven, K. J. and Kirkby, M. J. (1979). A physically-based, variable contributing area model of basin hydrology. *Hydrol. Sci. Bull.*, 24: 43–69.
- Beven, K. J., Lamb, R., Quinn, P., Romanowicz, R. and Freer, J. (1995). TOPMODEL. In: Singh, V. P. (Ed.), *Computer Models of Watershed Hydrology*. Water Resources Publications, Highlands Ranch, Colorado, pp.627–68.
- Beven, K. J. and Moore, I. D. (Eds) (1992). *Terrain Analysis and Distributed Modelling in Hydrology*. Wiley, New York, 249pp.

- Binley, A. M. and Beven, K. J. (1991). Physically-based modelling of catchment hydrology: a likelihood approach to reducing predictive uncertainty. In: Farmer, D. G. and Rycroft, M. J. (Eds), *Computer Modelling in the Environmental Sciences*, pp.75–88.
- Binley, A., Beven, K. and Elgy, J. (1989). A physically based model of heterogeneous hillslopes. 2. Effective hydraulic conductivities. *Water Resour. Res.*, 25: 1227–33.
- Blackman, R. B. and Tukey, J. W. (1958). *The Measurement of Power Spectra*. Dover Publ., New York, 190pp.
- Blöschl, G. (1996). *Scale and Scaling in Hydrology*. Institut für Hydraulik, Gewässerkunde und Wasserwirtschaft. Wiener Mitteilungen, Wasser-Abwasser-Gewässer, Vol. 132, Technical University of Vienna, 346pp.
- Blöschl, G. (1999). Scaling issues in snow hydrology. *Hydrological Processes*, 13, 2149–75.
- Blöschl, G., Grayson, R. B. and Sivapalan, M. (1995). On the representative elementary area (REA) concept and its utility for distributed rainfall-runoff modelling. *Hydrological Processes*, 9: 313–30.
- Blöschl, G., Gutknecht, D., Grayson, R. B., Sivapalan, M. and Moore, I. D. (1993). Organisation and randomness in catchments and the verification of distributed hydrologic models. *Eos, Transactions of the American Geophysical Union*, 74(43): 317.
- Blöschl, G., Gutknecht, D. and Kirnbauer, R. (1991a). Distributed snowmelt simulations in an alpine catchment. 2. Parameter study and model predictions. *Water Resour. Res.*, 27(12): 3181–8.
- Blöschl, G. and Kirnbauer, R. (1991). Point snowmelt models with different degrees of complexity – Internal processes. *J. Hydrology*, 129: 127–47.
- Blöschl, G. and Kirnbauer, R. (1992). An analysis of snow cover patterns in a small alpine catchment. *Hydrological Processes*, 6: 99–109.
- Blöschl, G., Kirnbauer, R. and Gutknecht, D. (1991b). Distributed snowmelt simulations in an Alpine catchment. 1. Model evaluation on the basis of snow cover patterns. *Water Resour. Res.*, 27: 3171–9.
- Blöschl, G. and Sivapalan, M. (1995). Scale issues in hydrological modelling – a review. *Hydrological Processes*, 9: 251–90.
- Blyth, E. M. and Harding, R. J. (1995). Application of aggregation models to surface heat flux from the Sahelian tiger brush. *Agric. For. Meteorol.*, 72: 213–35.
- Bonell, M. and Gilmour, D. A. (1978). The development of overland flow in a tropical rain-forest catchment. *J. Hydrology*, 39: 365–82.
- Bow, S.-T. (1992). *Pattern Recognition and Image Processing*. Dekker, New York, 558pp.
- Bowser, W. E., Peters, T. W. and Newton, J. D. (1951). Soil survey of the Red Deer sheet. *Alberta Soil Survey*, 16: 86pp.
- Bras, R. L. (1990). *Hydrology, An Introduction to Hydrologic Science*. Addison-Wesley, Reading, MA, 643pp.
- Braun, L. N. (1985). Simulation of snowmelt-runoff in lowland and lower alpine regions of Switzerland. Zürcher Geographische Schriften 21, Geographisches Institut der Eidgenössischen Technischen Hochschule Zürich.
- Bristow, K. L. and Campbell, G. S. (1984). On the relationship between incoming solar radiation and the daily maximum and minimum temperature. *Agric. For. Meteorol.*, 31: 159–66.
- Bronstert, A. and Plate, E. (1997). Modelling of runoff generation and soil moisture dynamics for hillslopes and micro-catchments. *J. Hydrology*, 198: 177–95.
- Brooks, R. H. and Corey, A. T. (1966). Properties of porous media affecting fluid flow. *J. Irrig. Drain. Div. Am. Soc. Civ. Eng.*, 92(IR2): 61–88.
- Brubaker, K. L. and Entekhabi, D. (1996). Analysis of feedback mechanisms in land-atmosphere interaction. *Water Resour. Res.*, 32(5): 1343–57.
- Brun, C., Bernard, R., Vidal-Madjar, D., Gascuel-Oudou, C., Mérot, P., Duchesne, J. and Nicolas, H. (1990). Mapping saturated areas with a helicopter-borne C band scatterometer. *Water Resour. Res.*, 26(5): 945–55.

- Brun, E., David, P., Sudul, M. and Brunot, G. (1992). A numerical model to simulate snow-cover stratigraphy for operational avalanche forecasting. *J. Glaciology*, 28(128): 13–22.
- Bruneau, P., Gascuel-Oudou, C., Robin, P., Mérot, P. and Beven, K. J. (1995). The sensitivity to space and time scales of a hydrological model using digital elevation data. *Hydrological Processes*, 9: 69–82.
- Brutsaert, W. (1982). *Evaporation into the Atmosphere*. Kluwer, Dordrecht, Holland, 299pp.
- Buol, S. W., Hole, F. D. and McCracken, R. J. (1989). *Soil Genesis and Classification*. Iowa State University Press: Ames.
- Burt, T. P. and Butcher, D. P. (1985). Topographic controls of soil moisture distributions. *J. Soil Science*, 36: 469–86.
- Burt, T. P. and Butcher, D. P. (1986). Development of topographic indices for use in semi-distributed hillslope runoff models. *Z. für Geomorph. N. F.*, Suppl. Bd. 58: 1–19.
- Calver, A. and Cammeraat, L. H. (1993). Testing a physically-based runoff model against field observations on a Luxembourg hillslope. *Catena*, 20(3): 273–88.
- Calver, A. and Wood, W. L. (1995). The Institute of Hydrology Distributed Model. In: Singh, V. P. (Ed.), *Computer Models of Watershed Hydrology*, Water Resources Publications, Highlands Ranch, Colorado, pp.595–626.
- Carlson, T. N., Gillies, R. R. and Perry, E. M. (1994). A method to make use of thermal infrared temperature and NDVI measurements to infer soil water content and fractional vegetation cover. *Remote Sens. Rev.*, 52: 45–59.
- Carlson, T. N., Perry, E. M. and Schmugge, T. J. (1990). Remote estimation of soil moisture availability and fractional vegetation cover for agricultural fields. *Agric. For. Meteorol.*, 52: 45–69.
- Carlson, V. A. (1969). Bedrock Topography of the Drumheller Map-Area, NTS 82P, Alberta, Research Council of Alberta.
- Charney, J. G., Halem, M. and Jastrow, R. (1969). Use of incomplete historical data to infer the present state of the atmosphere. *J. Atmos. Sci.*, 26: 1160–3.
- Chen, F. (1990a). Turbulent characteristics over a rough natural surface I: Turbulent structures. *Boundary-Layer Meteorol.*, 52: 151–75.
- Chen, F. (1990b). Turbulent characteristics over a rough natural surface II: Responses of profiles to turbulence. *Boundary-Layer Meteorol.*, 52, 301–11.
- Chow, V. T., Maidment, D. R. and Mays, L. W. (1988). *Applied Hydrology*. McGraw Hill.
- Cognard, A.-L., Loumagne, C., Normand, M., Olivier, P., Ottlé, C., Vidal-Madjar, D., Louahala, S. and Vidal, A. (1995). Evaluation of the ERS 1/synthetic aperture radar capacity to estimate surface soil moisture: Two-year results over the Naizin watershed. *Water Resour. Res.*, 31(4): 975–82.
- Colbeck, S. C. (1978). The physical aspects of water flow through snow. *Adv. Hydrosience*, 11: 165–206.
- Colbeck, S. C. (1988). Snowmelt increase through albedo reduction. Workshop on Snow Hydrology, November 1988, Manali (India), Central Water Commission and Himachal Pradesh State Electricity Board, India.
- Colbeck, S. C. (1991). The layered character of snow covers. *Rev. Geophys.*, 29(1): 81–96.
- Coles, N. A., Sivapalan, M., Larsen, J. E., Linnet, P. E. and Fahrner, C. K. (1997). Modelling runoff generation on small agricultural catchments: can real world runoff responses be captured? *Hydrological Processes*, 11: 111–36.
- Collier, C. G. and Knowles, J. M. (1986). Accuracy of rainfall estimates by radar III. Application for short-term flood forecasting. *J. Hydrology* 83: 237–49.
- Cooley, K. R. (1988). Snowpack variability on Western Rangelands. In: Western Snow Conference Proceedings, Kalispell, Montana, April 18–20.
- Cooley, R. L. (1997). Confidence intervals for ground-water models using linearization, likelihood and bootstrap methods. *Groundwater*, 35(5): 869–80.

- Coppin, P. A., Raupach, M. R. and Legg, B. J. (1986). Experiments on scalar dispersion within a model plant canopy Part II: An elevated plane source. *Boundary-Layer Meteorol.*, 35: 167–91.
- Coptý, N., Rubin, Y. and Mavko, G. (1993). Geophysical–hydrological identification of field permeabilities through Bayesian updating. *Water Resour. Res.*, 29: 2813–25.
- Costa-Cabral, M. C. and Burges, S. J. (1994). Digital elevation model networks (DEMON): A model of flow over hillslopes for computation of contributing and dispersal areas. *Water Resour. Res.*, 30: 1681–92.
- Crawford, N. H. and Linsley, R. S. (1966). Digital simulation in hydrology: The Stanford Watershed model IV. Technical Report No. 39. Dept. Civil Engineering, Stanford University, Palo Alto, California.
- Cressie, N. (1991). *Statistics for Spatial Data*. Wiley, New York, 900pp.
- Cunning, J. B. (1986). The Oklahoma–Kansas Preliminary Regional Experiment for STORM-Central, *Bull. Am. Meteor. Soc.*, 67(12): 1478–86.
- Daley, R. (1991). *Atmospheric Data Analysis*. Cambridge University Press, Cambridge, Massachusetts, 457pp.
- Danskin, W. R. (1988). Preliminary evaluation of the hydrogeologic system in Owens Valley, California. U.S.G.S. Water-Resources Investigations Report 88-4003.
- Davis, R. E., McKenzie, J. C. and Jordan, R. (1995). Distributed snow process modelling: an image processing approach. *Hydrological Processes*, 9: 865–75.
- Dawdy, D. R. (1969). Considerations involved in evaluating mathematical modelling of urban water systems. U.S. Geol. Survey Water Supply Paper 159I-D.
- Dawes, W. and Short, D. L. (1994). The significance of topology for modeling the surface hydrology of fluvial landscapes. *Water Resour. Res.*, 30(4): 1045–55.
- Dawes, W. R., Zhang, L., Hatton, T. J., Reece, P. H., Beale, G. T. H. and Packer, I. (1997). Evaluation of a distributed parameter ecohydrological model (Topog_IRM) on a small cropping rotation catchment. *J. Hydrology*, 191: 64–86.
- de Bruin, H. A. R., Bink, N. J. and Kroon, L. J. M. (1991). Fluxes in the surface layer under advective conditions. In: Schmugge, T. J. and André, J. C. (Eds), *Land Surface Evaporation Measurement and Parameterization*. Springer-Verlag, New York, pp.157–69.
- de Marsily, G. (1986). *Quantitative Hydrogeology*. Academic Press, San Diego, 440pp.
- de Marsily, G., Combes, P. and Goblet, P. (1992). Comments on ‘Ground-water models cannot be validated’, by L. F. Konikow and J. D. Bredehoeft. *Adv. Water Resour.*, 15: 367–9.
- De Troch, F. (1977). Studie van de oppervlaktewaterhydrologie van een Oost-Vlaams stroombekken: de Zwalmbeek te Nederzwalm. PhD dissertation, University of Gent, Faculty of Applied Sciences (in Dutch).
- De Troch, F. P., Troch, P. A., Su, Z. and Lin, D. S. (1996). Application of remote sensing for hydrological modelling. In: Abbott, M. B. and Refsgaard, J. C. (Eds), *Distributed Hydrological Modelling*. Kluwer, Dordrecht, The Netherlands, pp.165–91.
- Dee, D. P. (1995). A pragmatic approach to model validation. In: Lynch, D. R. and Davies, A. M. (Eds), *Quantitative Skill Assessment of Coastal Ocean Models*. AGU, Washington, pp.1–13.
- Deutsch, C. V. and Journel, A. G. (1997). *GSLIB, Geostatistical Software Library and User’s Guide*. Oxford University Press, New York, 384pp.
- Diak, G. R., Mecikalski, J. R., Anderson, M. C. and Bland, W. L. (1998). *Satellite Estimates of Terrestrial Radiation for Agricultural Applications*. 23rd Conference on Agricultural and Forest Meteorology, Albuquerque, NM. American Meteorological Society, Boston, MA, pp.1–2.
- Dickinson, R. E., Henderson-Sellers, A. and Kennedy, P. J. (1993). Biosphere–Atmosphere Transfer Scheme (BATS) Version 1e as coupled to the NCAR Community Climate Model. NCAR/TN-387+STR, National Center for Atmospheric Research.
- Dingman, S. L. (1994). *Physical Hydrology*, Macmillan, 575pp.

- Diskin, M. H. and Lane, L. J. (1972). A basinwide stochastic model for ephemeral stream runoff in South-eastern Arizona, *Bull. IASH*, 17(1): 61–76.
- Diskin, M. H. (1970). Definition and uses of the linear regression model. *Water Resour. Res.*, 6(6): 1668–73.
- Dolman, A. J. and Wallace, J. S. (1991). Lagrangian and *K*-theory approaches in modelling evaporation from sparse canopies. *Quart. J. R. Meteorol. Soc.*, 117: 1325–40.
- Dozier, J. (1979). A solar radiation model for a snow surface in mountainous terrain. In: Colbeck, S. C. and Ray, M. (Eds), *Modeling Snow Cover Runoff*. Proc. Conf. Hanover. U.S. Army CRREL, pp.144–53.
- Dozier, J. (1987). Recent research in snow hydrology. *Rev. Geophys.*, 25(2): 153–61.
- Dozier, J. and Frew, J. (1990). Rapid calculation of terrain parameters for radiation modelling from digital elevation data. *IEEE Transactions on Geoscience and Remote Sensing* 28(5): 963–9.
- Droegemeier, K. K. (1997). The numerical prediction of thunderstorms: challenges, potential benefits and results from realtime operational tests, *WMO Bull.*, 46: 324–36.
- Droegemeier, K. K., Xue, M., Sathye, A., Brewster, K., Bassett, G., Zhang, J., Liu, Y., Zou, M., Crook, A., Wong, V. and Carpenter, R. (1996a). *Realtime Numerical Prediction of Storm-Scale Weather during VORTEX '95, Part I: Goals and Methodology*. Preprints, 18th Conf. on Severe Local Storms, 15–20 Jan., Amer. Meteor. Soc., San Francisco, CA, pp. 6–10.
- Droegemeier, K. K., Xue, M., Brewster, K., Liu, Y., Park, S. K., Carr, F., Mewes, J., Zong, J., Sathye, A., Bassett, G., Zou, M., Carpenter, R., McCarthy, D., Andra, D., Janish, P., Graham, R., Sanielvici, S., Brown, J., Loftis, B. and McLain, K. (1996b). *The 1996 CAPS Spring Operational Forecasting Period – Realtime Storm-Scale NWP, Part I: Goals and Methodology*. Preprints, 11th Conf. on Num. Wea. Pred., 19–23 August, Norfolk, VA, Amer. Meteor. Soc., San Francisco, CA, pp. 294–6.
- Duan, Q., Sorooshian, S. and Gupta, V. (1992). Effective and efficient global optimization for conceptual rainfall-runoff models. *Water Resour. Res.*, 28(4): 1015–31.
- Dubayah, R., Dozier, J. and Davis, F. W. (1990). Topographic distribution of clear-sky radiation over the Konza Prairie, Kansas. *Water Resour. Res.*, 26(4): 679–90.
- Dunne, T. (1978). Field studies of hillslope flow processes. In: Kirkby, M. J. (Ed.), *Hillslope Hydrology*. Wiley, Chichester, pp.227–93.
- Dunne, T. and Black, R. D. (1970a). An experimental investigation of runoff production in permeable soils. *Water Resour. Res.*, 6(2): 478–90.
- Dunne, T. and Black, R. D. (1970b). Partial area contributions to storm runoff in a small New England watershed. *Water Resour. Res.*, 6(5): 1296–311.
- Dunne, T., Moore, T. R. and Taylor, C. H. (1975). Recognition and prediction of runoff-producing zones in humid regions. *Hydrol. Sci. Bull.*, 20: 305–27.
- Eagleson, P. S. (1978a). Climate soil and vegetation, 1, Introduction to water balance dynamics. *Water Resour. Res.*, 14(5): 705–12.
- Eagleson, P. S. (1978b). Climate soil and vegetation, 2, The distribution of annual precipitation derived from observed storm sequences. *Water Resour. Res.*, 14(5): 713–21.
- Eagleson, P. S. (1978c). Climate soil and vegetation, 3, A simplified model of soil moisture movement in the liquid phase. *Water Resour. Res.*, 14(5): 722–30.
- Eagleson, P. S. (1978d). Climate soil and vegetation, 4, The expected value of annual evapotranspiration. *Water Resour. Res.*, 14(5): 731–40.
- Eagleson, P. S. (1978e). Climate soil and vegetation, 5, A derived distribution of storm surface runoff. *Water Resour. Res.*, 14(5): 741–8.
- Eagleson, P. S. (1978f). Climate soil and vegetation, 6, Dynamics of the annual water balance. *Water Resour. Res.*, 14(5): 749–64.
- Eagleson, P. S., Fennessey, N. M., Qinling, W. and Rodriguez-Iturbe, I. (1987). Application of spatial Poisson models to air-mass thunderstorm rainfall. *J. Geophys. Res.*, 92(D8): 9661–78.

- Elder, K. (1995). Snow Distribution in Alpine Watersheds. Ph.D Thesis, Department of Geography, University of California.
- Elder, K., Michaelsen, J. and Dozier, J. (1995). Small basin modelling of snow water equivalence using binary regression tree methods. In: Tonnessen, K. A. et al. (Eds), *Biogeochemistry of Seasonally Snow-Covered Catchments*, Proc. Boulder Symposium, July 3–14. IAHS Publ. No. 228, pp.129–39.
- Elder, K., Rosenthal, W. and Davis, R. (1998). Estimating the spatial distribution of snow water equivalence in a montane watershed. *Hydrological Processes*, 12(10–11): 1793–808.
- Elsenbeer, H., Cassel, K. and Castro, J. (1992). Spatial analysis of soil hydraulic conductivity in a tropical rainforest catchment. *Water Resour. Res.*, 28(12): 3201–14.
- Elsenbeer, H., Cassel, K. and Zuniga, L. (1994). Throughfall in the terra firme forest of western Amazonia. *J. Hydrology (N.Z.)*, 32(2): 30–44.
- Elsenbeer, H. and Lack, A. (1996). Hydrometric and hydrochemical evidence for fast flow-paths at La Cuenca, Western Amazonia. *J. Hydrology*, 180: 237–50.
- Elsenbeer, H., Lack, A. and Cassel, K. (1995). Chemical fingerprints of hydrological compartments and flowpaths at La Cuenca, Western Amazonia. *Water Resour. Res.*, 31(12): 3051–8.
- Elsenbeer, H. and Vertessy, R. A. (2000). Stormflow generation and flowpath characteristics in an Amazonian rainforest catchment. *Hydrological Processes*, 14, 2367–2381.
- Emmett, W. W. (1970). The hydraulics of overland flow on hillslopes. U.S. Geol. Survey Prof. Paper 662-A, 47pp.
- Engman, E. T. (1986). Roughness coefficients for routing surface runoff. *J. Irrigation and Drainage Div., Proc ASCE*, 112: 39–53.
- Engman, E. T. (1991). Applications of microwave remote sensing of soil moisture for water resources and agriculture. *Remote Sens. Environ.*, 35: 213–26.
- Engman, E. T. (1995). Recent advances in remote sensing in hydrology. *Reviews of Geophysics, Supplement*, 967–75.
- Entekhabi, D. and Eagleson, P. S. (1989). Land surface hydrology parameterization for atmospheric general circulation models including subgrid scale spatial variability. *J. Climate*, 2: 816–31.
- Ericksen, B. and Myrabø, S. (1990). Studies of the relationship between soil moisture and topography in a small catchment. In: Gambolati, G. (Ed.), *Proc. 8th International Conference on Computational Methods in Water Resources*, Venice.
- Essery, R., Li, L. and Pomeroy, J. (1999). A distributed model of blowing snow over complex terrain. *Hydrological Processes*, 13: 2423–38.
- Ewen, G. O. and Parkin, G. (1996). Validation of catchment models for predicting land-use and climate change impacts. 1. Method. *J. Hydrology*, 175: 583–94.
- Famiglietti, J. S. and Wood, E. F. (1994). Multiscale modeling of spatially variable water and energy balance processes. *Water Resour. Res.*, 30: 3061–78.
- Famiglietti, J. S. and Wood, E. F. (1995). Effects of spatial variability and scale on areally averaged evapotranspiration. *Water Resour. Res.*, 31: 699–712.
- Fan, Y. and Bras, R. L. (1995). On the concept of a representative elementary area in catchment runoff. *Hydrological Processes*, 9: 821–32.
- Farajalla, N. M. and Vieux, B. E. (1995). Capturing the essential spatial variability in distributed hydrological modelling: infiltration parameters. *Hydrological Processes*, 9: 55–68.
- Faurès, J. M., Goodrich, D. C., Woolhiser, D. A. and Sorooshian, S. (1995). Impact of small-scale spatial rainfall variability on runoff modeling. *J. Hydrology*, 173: 309–26.
- Federico, V. D. and Neuman, S. P. (1997). Scaling of random fields by means of truncated power variograms and associated spectra. *Water Resour. Res.*, 33(5): 1075–85.
- Ferré, P. A., Knight, J. H., Rudolph, D. L. and Kachanoski, R. G. (1998). The sample areas of conventional and alternative time domain reflectometry probes. *Water Resour. Res.*, 34(11): 2971–9.

- Finn, D., Lamb, B., Leclerc, M. Y. and Horst, T. W. (1996). Experimental evaluation of analytical and lagrangian surface-layer flux footprint models. *Boundary-Layer Meteorol.*, 80: 282–308.
- Flavelle, P. (1992). A quantitative measure of model validation and its potential use for regulatory purposes. *Adv. Water Resour.*, 15: 5–13.
- Flury, M., Flühler, H., Jury, W. A. and Leuenberger, J. (1994). Susceptibility of soils to preferential flow of water: A field study. *Water Resour. Res.*, 30: 1945–54.
- Forkel, C. (1996). Das numerische Modell – ein schmaler Grat zwischen vertrauenswürdigem Werkzeug und gefährlichem Spielzeug. Presented at the 26th IWASA, RWTH Aachen, 4–5 January 1996.
- Foufoula-Georgiou, E. and Georgakakos, K. P. (1991). Recent advances in space-time precipitation modeling and forecasting. Chapter 3 in Bowles, D. and O’Connell, E. (Eds), *Recent Advances in the Modeling of Hydrological Systems*. Reidel Publ. Co., pp. 47–65.
- Foufoula-Georgiou, E. and Krajewski, W. (1995). Recent advances in rainfall modeling, estimation and forecasting. *Rev. Geophys. IUGG, U.S National Report, 1991–1994 – Contributions in Hydrology*.
- Foufoula-Georgiou, E. and Tsonis, T. (Eds) (1996). Space-time variability and dynamics of rainfall. *J. Geophys. Res.* (Special issue), 101(D21): 26161–538.
- Franks, S., Beven, K. J. and Lamb, R. (1997). Assessing the utility of distributed data in constraining the behaviour of hydrological models in the Wijlegemse subcatchment. In: De Troch, F. P., Troch, P. A. and Su, Z. (Eds), *Spatial and Temporal Soil Moisture Mapping from ERS-1 and JERS-1 SAR Data and Macroscale Hydrological Modelling*. Final Research Report, EC Environment Research Program EV5V-CT94-0446. Gent, Belgium, pp.334–57.
- Franks, S. W., Gineste, Ph., Beven, K. J. and Mérot, Ph. (1998). On constraining the predictions of a distributed model: The incorporation of fuzzy estimates of saturated areas into the calibration process. *Water Resour. Res.*, 34(4): 787–97.
- Freer, J., Beven, K. and Ambrose, B. (1996). Bayesian-estimation of uncertainty in runoff prediction and the value of data: An application of the GLUE approach. *Water Resour. Res.*, 32(7): 2161–73.
- Freeze, R. A. (1971). Three-dimensional, transient, saturated-unsaturated flow in a ground-water basin. *Water Resour. Res.*, 7(2): 347–66.
- Freeze, R. A. (1980). A stochastic-conceptual analysis of rainfall-runoff processes on a hill-slope. *Water Resour. Res.*, 16(2): 391–408.
- Freeze, R. A. and Harlan, R. L. (1969). Blueprint for a physically-based, digitally simulated hydrologic response model. *J. Hydrology*, 9: 237–58.
- Freeze, R. A. and Witherspoon, P. A. (1966). Theoretical analysis of regional groundwater flow: 1. Analytical and numerical solutions to the mathematical model. *Water Resour. Res.*, 2(4): 641–56.
- Frew, J. E. (1990). The Image Processing Workbench, PhD Thesis, University of California, Santa Barbara.
- Friedl, M. A. (1997). Examining the effects of sensor resolution and sub-pixel heterogeneity on spectral vegetation indices: Implications for biophysical modeling. In: Quattrochi, D. A. and Goodchild, M. F. (Eds), *Scale in Remote Sensing and GIS*. CRC Lewis Pub., New York, pp.113–39.
- Fritsch, J. M., Houze, R. A. Jr., Adler, R., Bluestein, H., Bosart, L., Brown, J., Carr, F., Davis, C., Johnson, R. H., Junker, N., Kuo, Y. H., Rutledge, S., Smith, J., Toth, Z., Wilson, J. W., Zipser, E. and Zrnich, D. (1998). Quantitative Precipitation Forecasting: Report of the Eight Prospectus Development Team, U.S. Weather Research Program, *Bull. Amer. Meteor. Soc.*
- Gardner, W. R. (1958). Some steady state solutions of the unsaturated moisture flow equation with application to evaporation from a water table. *Soil Science*, 85(4): 228–32.
- Garratt, J. R. (1978). Flux profile relations above tall vegetation. *Quart. J. R. Meteorol. Soc.*, 104: 199–211.

- Garratt, J. R. (1980). Surface influences upon vertical profiles in the atmospheric near-surface layer. *Quart. J. R. Meteorol. Soc.*, 106: 803–19.
- Garratt, M. J. and Spencer-Jones, D. (1981). *Yan Yean Geological Map*. Geological Survey of Victoria, Melbourne.
- Gelhar, L. W. (1993). *Stochastic Subsurface Hydrology*. Prentice Hall, Englewood Cliffs, NJ, 390pp.
- Gelman, A., Carlin, J. B., Stren, H. S. and Rubin, D. B. (1997). *Bayesian Data Analysis*. Chapman and Hall, p.526.
- General Meteorological Package (GEMPAK) (1992). Version 5.1, Reference manual, Unidata Program Center, UCAR, Boulder.
- Gessler, P. E, Moore, I. D., McKenzie, N. J. and Ryan, P. J. (1995). Soil landscape modelling and spatial prediction of soil attributes. *Int. J. Geogr. Inf. Sys.*, 9(4): 421–32.
- Ghan, S. J., Liljegren, J. C., Shaw, W. J., Hubbe, J. H. and Doran, J. C. (1997). Influence of subgrid variability on surface hydrology. *J. Climate*, 10: 3157–66.
- Gillies, R. R. and Carlson, T. N. (1995). Thermal remote sensing of surface soil water content with partial vegetation cover for incorporation into climate models. *J. Appl. Meteorol.*, 34: 745–56.
- Gillies, R. R., Cui, J., Carlson, T. N., Kustas, W. P. and Humes, K. S. (1997). Verification of a method for obtaining surface soil water content and energy fluxes from remote measurements of NDVI and surface radiant temperature. *Int. J. Remote Sens.*, 18: 3145–66.
- Gineste, P., Puech, C. and Mérot, P. (1998). Radar remote sensing of the source areas from the Coët-Dan catchment. *Hydrological Processes*, 12: 267–84.
- Giorgi, F. and Avissar, R. (1997). Representation of heterogeneity effects in earth system modeling: Experience from land surface modeling. *Rev. Geophys.*, 35: 413–38.
- Gomendy, V., Galhi, M., Western, A., Grayson, R. and White, R. (in preparation). Description and spatial variability of a yellow Duplex and Grey massive earth soil toposequence at the small catchment scale in Victoria.
- Goodrich, D. C. (1990). Geometric simplification of a distributed rainfall-runoff model over a range of basin scales. PhD Thesis, The University of Arizona, 361pp.
- Goodrich, D. C., Faurés, J.-M., Woolhiser, D. A., Lane, L. J. and Sorooshian, S. (1995). Measurement and analysis of small-scale convective storm rainfall variability. *J. Hydrology*, 173: 283–308.
- Goodrich, D. C., Lane, L. J., Shillito, R. A., Miller, S. N., Syed, K. H. and Woolhiser, D. A. (1997). Linearity of basin response as a function of scale in a semi-arid watershed. *Water Resour. Res.* 33(12): 2951–65.
- Goodrich, D. C., Schumge, T. J., Jackson, T. J., Unkrich, C. L., Keefer, T. O., Parry, R., Bach, L. B. and Amer, S. A. (1994). Runoff simulation sensitivity to remotely sensed initial soil water content. *Water Resour. Res.*, 30(5): 1393–405.
- Goodrich, D. C., Stone, J. J. and Van der Zweep, R. (1993). Validation Strategies Based on Model Application Objectives. Proc. Fed. Interagency Workshop on Hydrologic Modelling Demands for the 90's, U.S.G.S. Water Resources Invest. Report 93-4018, Ft. Collins, CO, June 7-9, pp.8-1–8-8.
- Grayson, R. B., Blöschl, G., Barling, R. D. and Moore, I. D. (1993). Process, scale and constraints to hydrological modelling in GIS. In: Kovar, K. and Nachtnebel, H. P. (Eds), *Applications of Geographic Information Systems in Hydrology and Water Resources Management*. Proc. Vienna Symp., April 1993, IAHS Publ. No. 211, pp. 83–92.
- Grayson, R. B., Blöschl, G. and Moore, I. D. (1995). Distributed parameter hydrologic modelling using vector elevation data: THALES and TAPES-C. Chapter 19 in: Singh, V. P. (Ed.), *Computer Models of Watershed Hydrology*. Water Resources Publications, Highlands Ranch, Colorado, pp.669–96.
- Grayson, R. B., Moore, I. D. and McMahon, T. A. (1992a). Physically based hydrologic modeling, 1. A terrain-based model for investigative purposes. *Water Resour. Res.*, 28: 2639–58.

- Grayson, R. B., Moore, I. D. and McMahon, T. A. (1992b). Physically based hydrologic modeling, 2. Is the concept realistic? *Water Resour. Res.*, 28: 2659–66.
- Grayson, R. B. and Western, A. W. (1998). Towards areal estimation of soil water content from point measurements: time and space stability of mean response. *J. Hydrology*, 207: 68–82.
- Grayson, R. B., Western, A. W., Chiew, F. H. S. and Blöschl, G. (1997). Preferred states in spatial soil moisture patterns: Local and nonlocal controls. *Water Resour. Res.*, 33(12): 2897–908.
- Gupta, V. K. and Waymire, E. (1979). A stochastic kinematic study of subsynoptic space-time rainfall. *Water Resour. Res.*, 15(3): 637–44.
- Gupta, V. K. and Waymire, E. (1990). Multiscaling properties of spatial rainfall in river flow distributions. *J. Geophys. Res.*, 95: 1999–2009.
- Gupta, H. V., Sorooshian, S. and Yapo, P. O. (1998). Toward improved calibration of hydrologic models: Multiple and noncommensurable measures of information. *Water Resour. Res.*, 34(4): 751–63.
- Gyasi-Agyei, Y., Willgoose, G. and De Troch, F. P. (1995). Effects of vertical resolution and map scale of digital elevation models on geomorphologic parameters used in hydrology. *Hydrological Processes*, 9: 363–82.
- Hardy, J., Albert, M. and Marsh, P. (Eds) (1999). Snow hydrology, *Hydrological Processes* (special issue), 13: Nos 12–15.
- Hatton, T. J., Dawes, W. R. and Vertessy, R. A. (1995). The importance of landscape position in scaling SVAT models to catchment scale hydroecological prediction. In: Feddes, R. A. (Ed.), *Space and Time Scale Variability and Interdependencies in Hydrological Processes*. Cambridge University Press, New York, pp.43–53.
- Henebry, G. M. (1997). Advantages of principal components analysis for land cover segmentation from SAR image series. In: *Proceedings of the 3rd ERS symposium*. Florence, Italy, pp.175–8.
- Herschy, R. W. (Ed.) (1999). *Hydrometry: Principles and Practice*. Wiley, 384pp.
- Hewlett, J. D. and Hibbert, A. R. (1967). Factors affecting the response of small watersheds to precipitation in humid areas. In: Sopper, W. E. and Lull, H. W. (Eds), *Forest Hydrology*. Pergamon Press, Oxford, pp.275–90.
- Hillel, D. (1986). Modelling in soil physics: A critical review. In: *Future Developments in Soil Science Research*. A collection of Soil Sci. Soc. Am. Golden Anniversary contributions presented at Annual Meeting, New Orleans, pp.35–42.
- Hipps, L. E., Humes, K. and Kustas, W. P. (1995). *Defining Scales of Inhomogeneity for Surface Properties: Implications towards Studies of Surface–Atmosphere Interactions*. Proceedings of Conference on Hydrology, Dallas, TX, American Meteorological Society, Boston, MA, pp.141–2.
- Hipps, L. E., Or, D. and Neale, C. M. U. (1996). Spatial structure and scaling of surface fluxes in a Great Basin ecosystem. In: Stewart, J. B., Engman, E. T., Feddes, R. A. and Kerr, Y. (Eds), *Scaling up in Hydrology using Remote Sensing*. Wiley, London, pp.113–25.
- Hipps, L. E., Swiatek, E. and Kustas, W. P. (1994). Interactions between regional surface fluxes and the atmospheric boundary layer over a heterogeneous watershed. *Water Resour. Res.*, 30(5): 1387–92.
- Hochstöger, F. (1989). Ein Beitrag zur Anwendung und Visualisierung digitaler Geländemodelle (On the application and visualization of digital terrain models). *Geowissenschaftliche Mitteilungen* 34, Veröffentlichungen des Institutes für Photogrammetrie und Fernerkundung, Technische Universität Wien.
- Hornberger, G. M. (1994). Data and analysis note: A new type of article for Water Resources Research. *Water Resour. Res.*, 30(12): 3241–2.
- Horst, T. W. and Weil, J. C. (1992). Footprint estimation for scalar flux measurements in the atmospheric surface layer. *Boundary-Layer Meteorol.*, 59: 279–96.

- Horton, R. E. (1933). The role of infiltration in the hydrologic cycle. *Trans. Am. Geophys. Union*, 14: 446–60.
- Houser, P. R. (1996). Remote Sensing Soil Moisture using 4-Dimensional Data Assimilation. Ph.D dissertation, Department of Hydrology and Water Resources, The University of Arizona.
- Houser, P. R., Gupta, H. V., Shuttleworth, W. J. and Famiglietti, J. S. (2000). Multi-objective calibration and sensitivity of a distributed land–surface water and energy balance model. *Water Resour. Res.* (in press)
- Houser, P. R., Shuttleworth, W. J., Gupta, H. V., Famiglietti, J., Syed, K. H. and Goodrich, D. C. (1998). Integration of soil moisture remote sensing and hydrologic modelling using data assimilation. *Water Resour. Res.*, 34(12): 3405–20.
- Huff, F. A. (1967). Time distribution of rainfall in heavy storms. *Water Resour. Res.*, 3(4): 1007–19.
- Humes, K. S., Kustas, W. P. and Goodrich, D. C. (1997). Spatially distributed sensible heat flux over a semiarid watershed. Part I: use of radiometric surface temperatures and spatially uniform resistance. *J. Appl. Meteorol.*, 36: 281–92.
- Huntingford, C., Allen, S. J. and Harding, R. J. (1995). An intercomparison of single and dual-source vegetation–atmosphere transfer model applied to transpiration from sahelian savannah. *Boundary-Layer Meteorol.*, 74: 397–418.
- Hursh, C. R. and Brater, E. F. (1941). Separating storm-hydrographs from small drainage-areas into surface- and subsurface-flow. *Trans. American Geophysical Union*, 22: 863–70.
- Hutchinson, M. F. (1989). A new method for gridding elevation and stream line data with automatic removal of spurious pits. *J. Hydrology*, 106: 211–32.
- Hutchinson, M. F. (1991). The application of thin plate smoothing splines to continent-wide data assimilation. In: Jasper, J. D. (Ed.), *Data Assimilation Systems*, BMRC Research Report 27, Bureau of Meteorology, Melbourne, 104–13.
- Hutchinson, M. F. (1993). On thin plate splines and kriging. In: Tarter, M. E. and Lock, M. D. (Eds), *Computing Science and Statistics*, Vol.25. Interface Foundation of North America, University of California, Berkeley, pp.55–62.
- Hutchinson, M. F. and Gessler, P. E. (1994). Splines – more than just a smooth interpolator. *Geoderma*, 62: 45–67.
- IAHR (1994). Publication of guidelines for validation documents and call for discussion. *International Association for Hydraulic Research Bulletin*, 11:41.
- Iorgulescu, I. and Musy, A. (1997). Generalisation of TOPMODEL for a power law transmissivity profile. *Hydrological Processes*, 11: 1353–5.
- Islam, S., Bras, R. L. and Rodriguez-Iturbe, I. (1988). Multidimensional modelling of cumulative rainfall: Parameter estimation and model adequacy through a continuum of scales. *Water Resour. Res.*, 24(7): 985–92.
- Jackson, T. H. R. (1994). A Spatially Distributed Snowmelt-Driven Hydrologic Model applied to the Upper Sheep Creek Watershed. Ph.D Thesis, Utah State University.
- Jackson, T. J. and Le Vine, D. E. (1996). Mapping surface soil moisture using an aircraft-based passive microwave instrument: algorithm and example. *J. Hydrology*, 184: 57–84.
- Jackson, T. J., Le Vine, D. M., Griffis, A. J., Goodrich, D. C., Schmutge, T. J., Swift, C. T. and O'Neill, P. E. (1993). Soil moisture and rainfall estimation over a semiarid environment with the ESTAR microwave radiometer. *IEEE Trans. Geosci. Remote Sens.*, 31(4): 836–41.
- Jacobs, B. L., Rodriguez-Iturbe, I. and Eagleson, P. S. (1988). Evaluation of a homogeneous point process description of Arizona thunderstorm rainfall. *Water Resour. Res.*, 24(7): 1174–86.
- Jain, S. K., Storm, B., Bathurst, J. C., Refsgaard, J. C. and Singh, R. D. (1992). Application of the SHE to catchments in India. Part 2. Field experiments and simulation

- studies with the SHE on the Kolar subcatchment of the Narmada River. *J. Hydrology*, 140: 25–47.
- Jakeman, A. and Hornberger, G. M. (1993). How much complexity is warranted in a rainfall–runoff model? *Water Resour. Res.*, 29(8): 2637–49.
- James, L. D. and Burges, S. J. (1982). Selection, calibration and testing of hydrological models. In: Haan, C. T., Johnson, H. P. and Brakensiek, D. L. (Eds), *Hydrological Modelling of Small Watersheds*. Am. Soc. Agr. Eng. Monogr. Series Vol. 5, pp.435–72.
- Jarvis, P. G. and McNaughton, K. G. (1986). Stomatal control of transpiration: Scaling up from leaf to region. *Adv. Ecological Research*, 15: 1–49.
- Jayatilaka, C. J., Storm, B. and Mudgway, L. B. (1998). Simulation of water flow on irrigation bay scale with MIKE SHE, *J. Hydrology*, 208: 108–30.
- Jensen, H. (1989). Räumliche Interpolation der Stundenwerte von Niederschlag, Temperatur und Schneehöhe. Zürcher Geographische Schriften 35. Geographisches Institut der Eidgenössischen Technischen Hochschule Zürich, 70pp.
- Jensen, K. H. and Mantoglou, A. (1993). Future of distributed modelling. Chapter 12 in: Beven, K. J. and Moore, I. D. (Eds), *Terrain Analysis and Distributed Modelling in Hydrology*. Wiley, New York, pp.203–12.
- Jones, E. B. (1983). Snowpack Ground-Truth Manual, Report No. CR 170584, National Aeronautics and Space Administration (NASA), Goddard Space Flight Center, Greenbelt, MD.
- Jordan, J. P. (1994). Spatial and temporal variability of stormflow generation processes on a Swiss catchment. *J. Hydrology*, 153: 357–82.
- Jordan, R. (1991). A one-dimensional temperature model for a snow cover. Technical documentation for SNTherm.89, special technical report 91–16, U.S. Army CRREL.
- Journel, A. G. and Deutsch, C. V. (1993). Entropy and spatial disorder. *Math. Geol.*, 25: 329–55.
- Journel, A. G. and Huijbregts, C. J. (1978). *Mining Geostatistics*, Academic Press, London, 600pp.
- Kabat, P., Hutjes, R. W. A. and Feddes, R. A. (1997). The scaling characteristics of soil parameters: From plot scale heterogeneity to subgrid parameterization. *J. Hydrology*, 190: 363–96.
- Kader, B. A. and Yaglom, A. M. (1990). Mean fields and fluctuation moments in unstably stratified turbulent boundary layers. *J. Fluid Mech.*, 212: 637–62.
- Kain, J. S. and Fritsch, J. M. (1992). The role of convective “trigger function” in numerical weather forecasts of mesoscale convective systems. *Meteor. Atmos. Phys.*, 40: 93–106.
- Karnieli, A. M., Diskin, M. H. and Lane, L. J. (1994). CELMOD5 – a semi-distributed cell model for conversion of rainfall into runoff in semi-arid watersheds. *J. Hydrology*, 157: 61–85.
- Katul, G., Goltz, S. M., Hsieh, C., Cheng, Y., Mowry, F. and Sigmon, J. (1995). Estimation of surface heat and momentum fluxes using the flux-variance method above uniform and non-uniform terrain. *Boundary-Layer Meteorol.*, 74: 237–60.
- Kavvas, M. L. and Delleur, J. W. (1981). A stochastic cluster model for daily rainfall sequences. *Water Resour. Res.*, 17(4): 1151–60.
- Keller, R. (Ed.) (1978). *Hydrologischer Atlas der Bundesrepublik Deutschland*, Deutsche Forschungsgemeinschaft. Boldt Verlag, Boppard, Germany, 68pp.
- Kendall, C. and McDonnell, J. J. (Eds) (1998). *Isotope Tracers in Catchment Hydrology*. Elsevier, 839pp.
- Khanna, S. and Brasseur, J. G. (1997). Analysis of Monin–Obukhov similarity from large-eddy simulation. *J. Fluid Mech.*, 345: 251–86.
- Kiefer, C. J. and Chu, H. H. (1957). Synthetic storm pattern for drainage design. *J. Hydr. Div., ASCE*, 83(HY4): 1–25.

- Kim, C. P., Salvucci, G. D. and Entekhabi, D. (1999). Groundwater–surface water interaction and the climatic spatial patterns of hillslope hydrological response. *Hydrology and Earth Systems Science*, 3(3): 375–84.
- Kirkby, M. (1975). Hydrograph modelling strategies. In: Peel, R., Chisholm, M. and Haggett, P. (Eds), *Processes in Physical and Human Geography*, pp.69–90.
- Kirkby, M. J. (1988). Hillslope runoff processes and models. *J. Hydrology*, 100: 315–40.
- Kirkby, M. J. (1997). TOPMODEL: a personal view. *Hydrological Processes*, 11(9): 1087–97.
- Kirkby, M. J., Callan, J., Weyman, D. and Wood, J. (1976). Measurement and modeling of dynamic contributing areas in a very small catchment. University of Leeds, School of Geography, Working Paper 167, Leeds, UK.
- Kirnbauer, R. and Blöschl, G. (1990). A lysimetric snow pillow station at Kühtai/Tyrol. In: Lang, H. and Musy, A. (Eds), *Hydrology in Mountainous Regions. I – Hydrological Measurements; the Water Cycle*. Proc. Lausanne Symp., August 1990, IAHS Publ. no. 193, pp.173–80.
- Kirnbauer, R. and Blöschl, G. (1994). Wie ähnlich sind Ausaperungsmuster von Jahr zu Jahr? (How similar are snow cover patterns from year to year?) *Deutsche Gewässerkundliche Mitteilungen*, 37(5/6): 113–21.
- Kirnbauer, R., Blöschl, G. and Gutknecht, D. (1994). Entering the era of distributed snow models. *Nordic Hydrology*, 25: 1–24.
- Klemeš, V. (1983). Conceptualisation and scale in hydrology. *J. Hydrology*, 65: 1–23.
- Klemeš, V. (1986a). Dilettantism in hydrology: transition or destiny? *Water Resour. Res.*, 22: 177S–88S.
- Klemeš, V. (1986b). Operational testing of hydrological simulation models. *Hydrol. Sci. J.*, 31(1–3): 13–24.
- Kneizys, F. X., Shettles, E. P., Abreu, L. W., Chetwynd, J. H., Anderson, G. P., Anderson, W. O., Selby, J. E. A. and Clough, S. A. (1988). Users Guide to LOWTRAN 7, AFGL-TR-0177, Air Force Geophysical Laboratory.
- Knisel, W. G. (Ed.) (1980). *CREAMS: A field-scale model for chemical runoff and erosion from agricultural management system*. USDA-SEA Conservation Research Report No. 26, 643pp.
- Koltermann, C. E. and Gorelick, S. M. (1996). Heterogeneity in sedimentary deposits: A review of structure-imitating, process-imitating, and descriptive approaches. *Water Resour. Res.*, 32: 2617–58.
- König, M. and Sturm, M. (1998). Mapping snow distribution in the Alaskan Arctic using aerial photography and topographic relationships. *Water Resour. Res.*, 34(12): 3471–83.
- Konikow, L. F. (1978). Calibration of groundwater models. In: *Verification of Mathematical and Physical Models in Hydraulic Engineering*. American Society of Civil Engineering, New York, pp.87–93.
- Konikow, L. F. and Bredehoeft, J. D. (1992). Ground-water models cannot be validated. *Adv. Water Resour.*, 15: 75–83.
- Koterba, M. T. (1986). Differential influences of storm and watershed characteristics on runoff from ephemeral streams in southeastern Arizona. Ph.D dissertation, Department of Hydrology and Water Resources, University of Arizona, Tucson, Arizona.
- Kottogoda, N. T. and Rosso, R. (1996). *Introductory Statistical, Probability and Reliability Methods for Civil and Environmental Engineers*. McGraw-Hill, New York.
- Kouwen, N. and Garland, G. (1989). Resolution considerations in using radar rainfall data for flood forecasting. *Can. J. Civ. Eng.*, 16: 279–89.
- Krajewski, W. F., Venkataraman, L., Georgakakos, K. P. and Jain, S. C. (1991). A Monte Carlo study of rainfall sampling effect on a distributed catchment model. *Water Resour. Res.*, 27(1): 119–28.
- Kraus, K. and Blöschl, G. (1998). Fernerkundung zur Sicherung der Wiener Wasservorräte (Remote sensing for safeguarding the Viennese water supply). In: Ricica, K. and Haslinger, U. (Eds). *Wasserspuren – Nachhaltige Zukunftspfade*, MA22 Umweltschutz, Wien, pp.26–9.

- Krige, D. G. (1951). A statistical approach to some basic mine evaluation problems on the Witwatersrand. *J. Chem. and Metall. Soc. of South Africa*, 52: 119–39.
- Kuczera, G. and Parent, E. (1998). Monte Carlo assessment of parameter uncertainty in conceptual catchment models: The Metropolis Algorithm. *J. Hydrology*, 211(1–4): 69–85.
- Kumar, P. and Foufoula-Georgiou, E. (1993a). A multicomponent decomposition of spatial rainfall fields: 1. Segregation of large and small-scale features using wavelet transforms. *Water Resour. Res.*, 29(8): 2515–32.
- Kumar, P. and Foufoula-Georgiou, E. (1993b). A multicomponent decomposition of spatial rainfall fields: 2. Self-similarity in fluctuations. *Water Resour. Res.*, 29(8): 2533–44.
- Kupfersberger, H. and Blöschl, G. (1995). Estimating aquifer transmissivities – on the value of auxiliary data. *J. Hydrology*, 165: 85–99.
- Kustas, W. P. and Daughtry, C. S. T. (1990). Estimation of the soil heat flux/net radiation ratio from multispectral data. *Agric. For. Meteorol.*, 49: 205–23.
- Kustas, W. P., Daughtry, C. S. T. and van Oevelen, P. J. (1993). Analytical treatment of the relationships between soil heat flux/net radiation ratio and vegetation indices. *Remote Sens. Environ.*, 46: 319–30.
- Kustas, W. P. and Goodrich, D. C. (1994). Preface to special section on Monsoon '90. *Water Resour. Res.*, 30(5): 1211–25.
- Kustas, W. P., Hips, L. E. and Humes, K. S. (1995). Calculation of basin-scale surface fluxes by combining remotely sensed data and atmospheric properties in a semiarid landscape. *Boundary-Layer Meteorol.*, 73: 105–24.
- Kustas, W. P. and Humes, K. S. (1996). Variations in the surface energy balance for a semi-arid rangeland using remotely sensed data at different spatial resolutions. In: Stewart, J. B., Engman, E. T., Feddes, R. A. and Kerr, Y. (Eds), *Scaling up in Hydrology using Remote Sensing.*, Wiley, London, pp.127–45.
- Kustas, W. P. and Norman, J. M. (1996). Use of remote sensing for evapotranspiration monitoring over land surfaces. *Hydrol. Sci. J.*, 41: 495–516.
- Kustas, W. P., Prueger, J. H., Hips, L. E., Ramalingam, K., Hatfield, J. L., Schmutge, T. J., Rango, A., Ritchie, J. C. and Havstad, K. M. (1998). *Application of Monin-Obukhov similarity over a mesquite dune site in the Jornada Experimental Range*. Proceedings of 23rd Conference on Agric. & Forest Meteorol., American Meteorological Society, Boston, MA, pp.216–19.
- Kustas, W. P., Rango, A. and Uijlenhoet, R. (1994). A simple energy budget algorithm for the snowmelt runoff model. *Water Resour. Res.*, 30(5): 1515–27.
- Kutchment, L. S., Demidov, V. N., Naden, P. S., Cooper, D. M. and Broadhurst, P. (1996). Rainfall-runoff modelling of the Oure basin, North Yorkshire: an application of a physically based distributed model. *J. Hydrology*, 181: 323–42.
- Lamb, R., Beven, K. and Myrabø, S. (1997). Discharge and water table predictions using a generalized TOPMODEL formulation. *Hydrological Processes*, 11(9): 1145–67.
- Lamb, R., Beven, K. and Myrabø, S. (1998a). A generalised topographic-soils hydrological index. In: Lane, S. N., Richards, K. S. and Chandler, J. H. (Eds), *Landform Monitoring, Modelling and Analysis*. Wiley, Chichester, pp.263–78.
- Lamb, R., Beven, K. and Myrabø, S. (1998b). Use of spatially distributed water table observations to constrain uncertainty in a rainfall-runoff model. *Adv. Water Resour.*, 22(4): 305–17.
- Lane, L. J. (1982). Distributed model for small semiarid watersheds. *J. Hydraulics Div., ASCE*, 108(HY10): 1114–31.
- Lane, L. J. (1983a). Transmission losses. Chapter 19 in: *SCS National Engr. Handbook*. US Government Printing Office, Washington, DC, pp.19-1–19-21.
- Lane, L. J. (1983b). *SPUR Hydrology Component: Water routing and sedimentation*, USDA-ARS Misc Pub. No. 1431, pp.62–7.
- Lane, L. J. and Renard, K. G. (1972). Evaluation of a basin wide stochastic model for ephemeral runoff from semiarid watersheds. *Trans. ASAE*, 15(1): 280–3.

- Laur, H., Bally, P., Meadows, P., Sanchez, J., Schaettler, B. and Lopinto, E. (1997). ERS SAR calibration: derivation of backscattering coefficient σ^0 in ESA ERS SAR PRI products (Tech. Rep.). Frascati, Italy: European Space Agency.
- Leavesley, G. H. and Stannard, L. G. (1995). The precipitation-runoff modelling system – PRMS. Chapter 9 in: Singh, V. P. (Ed.), *Computer Models of Watershed Hydrology*. Water Resources Publications, Highlands Ranch, Colorado, pp.281–310.
- Leclerc, M. Y. and Thurtell, G. W. (1990). Footprint prediction of scalar fluxes using a Markovian analysis. *Boundary-Layer Meteorol.*, 52: 247–58.
- Lee, J. and Hoppel, K. (1992). Principal components transformation of multifrequency polarimetric SAR imagery. *IEEE Trans. Geosc. Rem. Sens.*, 80(4): 686–96.
- Levine, J. B. and Salvucci, G. D. (1999a). Equilibrium analysis of groundwater–vadose zone interactions and the resulting spatial distribution of hydrologic fluxes across a Canadian prairie. *Water Resour. Res.*, 35(5): 1369–83.
- Levine, J. B. and Salvucci, G. D. (1999b). Characteristic rate scale and time scale of supply-limited transpiration under a Richards–Cowan framework. *Water Resour. Res.*, 35(12): 3947–54.
- Lhomme, J. P., Chehbouni, A. and Monteny, B. (1994). Effective parameters of surface energy balance in heterogeneous landscape. *Boundary Layer Meteorol.*, 71: 297–309.
- Lhomme, J. P., Monteny, B. and Bessemoulin, P. (1997). Inferring regional surface fluxes from convective boundary layer characteristics in a Sahelian environment. *Water Resour. Res.*, 33: 2563–9.
- Linsley, Jr., R. K., Kohler, M. A. and Paulhus, J. L. H. (1982). *Hydrology for Engineers*. McGraw-Hill.
- Liston, G. E. and Sturm, M. (1998). A snow-transport model for complex terrain. *J. Glaciology* 44: 498–516.
- Loague, K. and Kyriakidis, P. C. (1997). Spatial and temporal variability in the R-5 infiltration data set: Déjà vu and rainfall-runoff simulations. *Water Resour. Res.*, 33(12): 2883–95.
- Lopez, A., Laur, H. and Nezry, E. (1990). Statistical distribution and texture in multilook and complex SAR images. In: *Proc. International Geoscience and Remote Sensing Symposium (IGARSS)*. Washington DC., USA, pp.2427–30.
- Lopes, V. L., Nearing, M. A., Foster, G. R., Finkner, S. C. and Gilley, J. E. (1989). The Water Erosion Prediction Project: Model Overview. Proc. ASCE National Water Conf., Irrigation and Drain. and Water Resources Planning and Mgmt. Divs., Newark, DE, pp.503–10.
- Lørup, J. K., Refsgaard, J. C. and Mazvimavi, D. (1998). Assessing the effect of land use change on catchment runoff by combined use of statistical tests and hydrological modelling: Case studies from Zimbabwe. *J. Hydrology*, 205: 147–63.
- Los, H. and Gerritsen, H. (1995). Validation of water quality and ecological models. Presented at the 26th IAHR Conference, London, 11–15 Sept. 1995, Delft Hydraulics, 8pp.
- Luce, C. H., Tarboton, D. G. and Cooley, K. R. (1997). Spatially distributed snowmelt inputs to a semi-arid mountain watershed. Proc. Western Snow Conference, Banff, Canada, May 5–8.
- Luce, C. H., Tarboton, D. G. and Cooley, K. R. (1998). The influence of the spatial distribution of snow on basin-averaged snowmelt. *Hydrological Processes*, 12(10–11): 1671–83.
- Luce, C. H., Tarboton, D. G. and Cooley, K. R. (1999). Sub-grid parameterization of snow distribution for an energy and mass balance snow cover model. *Hydrological Processes*, 13: 1921–33.
- McClung, D. and Schaerer, P. (1993). *The Avalanche Handbook*, 2nd Edition. Mountaineers Books, 271pp.
- McDonald, M. G. and Harbaugh, A. W. (1996). User's documentation for MODFLOW-96, an update to the U.S. Geological Survey Modular Finite-Difference Ground-Water Flow Model, U.S. Geological Survey Open-File Report 96-485, 56pp.

- McDonnell, J. J., Bonell, M., Stewart, M. K. and Pearce, A. J. (1990). Deuterium variations in storm rainfall: implications for stream hydrograph separation. *Water Resour. Res.*, 26(3): 455–8.
- McKay, G. A. and Gray, D. M. (1981). The distribution of snowcover. Chapter 5 in: Gray, D. M. and Male, D. H. (Eds), *Handbook of Snow, Principles, Processes, Management and Use*. Pergamon Press, Willowdale, Canada.
- McLaughlin, D. and Townley, L. R. (1996). A reassessment of the groundwater inverse problem. *Water Resour. Res.*, 32(4): 1131–61.
- McNaughton, K. G. and Jarvis, P. G. (1983). Predicting effects of vegetation changes on transpiration and evaporation. In: Kozlowski, T. T. (Ed.), *Water Deficit and Plant Growth*, Vol. VII. Academic Press, New York, pp.2–47.
- McNaughton, K. G. and Spriggs, T. W. (1986). A mixed-layer model for regional evaporation. *Boundary Layer Meteorol.*, 34: 243–62.
- McNaughton, K. G. and Spriggs, T. W. (1989). An evaluation of the Priestley and Taylor equation and the complementary relationship using results from a mixed-layer model of the convective boundary layer. In: Black, T. A., Spittlehouse, D. L., Novak, M. D. and Price, D. T. (Eds), *Estimation of Areal Evapotranspiration*. IAHS Publ. no 177, Wallingford, pp.89–104.
- McNaughton, K. G. and Van den Hurk, B. J. J. M. (1995). A ‘Lagrangian’ revision of the resistors in the two-layer model for calculating the energy budget of a plant canopy. *Boundary-Layer Meteorol.*, 74: 262–88.
- Mahrt, L. (1998). Flux sampling errors for aircraft and towers. *J. Atmospheric and Oceanic Technology*, 15: 416–29.
- Male, D. H. and Gray, D. M. (1981). Snowcover ablation and runoff. Chapter 9 in: Gray D. M. and Male, D. H. (Eds), *Handbook of Snow, Principles, Processes, Management and Use*. Pergamon Press, Willowdale, Canada, pp.360–436.
- Mallat, S. (1989). A theory for multiresolution signal decomposition: the wavelet representation. *IEEE Trans. on Pattern Anal. and Mach. Intel.*, 11(7): 674–93.
- Mantoglou, A. and Wilson, J. L. (1981). Simulation of random fields with the turning bands method. Report No. 264, Department of Civil Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts, 199pp.
- Marks, D. and Dozier, J. (1992). Climate and energy exchange at the snow surface in the alpine region of the Sierra Nevada, 2. Snow cover energy balance. *Water Resour. Res.*, 28(11): 3043–54.
- Marsan, D., Schertzer, D. and Lovejoy, S. (1996). Causal space-time multifractal processes: Predictability and forecasting of rain fields. *J. Geophys. Res.*, 101(D21): 26333–46.
- Marshall, S. E. and Warren, S. G. (1987). Parameterization of snow albedo for climate models. In: Goodison, R. E., Barry, R. G. and Dozier, J. (Eds) *Large Scale Effects of Seasonal Snowcover*, Proc. Vancouver Symposium, August 1987, IAHS Publ. No. 166, pp. 43–50.
- Matheron, G. (1965). *Les variables régionalisées et leur estimation*. Masson, Paris. Cited in de Marsily (1986).
- Matheron, G. (1973). The intrinsic random functions and their applications. *Adv. in Appl. Prob.*, 5: 438–68.
- Mattikalli, N. M., Engman, E. T., Jackson, T. J. and Ahuja, L. R. (1998). Microwave remote sensing of temporal variations of brightness temperature and near-surface soil water content during a watershed-scale field experiment, and its application to the estimation of soil physical properties. *Water Resour. Res.*, 34(9): 2289–99.
- Mecikalski, J. R., Diak, G. R., Anderson, M. C. and Norman, J. M. (1999). Estimating fluxes on continental scales using remotely-sensed data in an atmospheric–land exchange model. *J. Appl. Meteorol.*, 38(9): 1352–69.
- Meijerink, A. M. J., Brouwer, H. A. M., Mannaerts, C. M. and Valenzuela, C. R. (1994). Introduction to the use of geographic information systems for practical hydrology. Report No. 23, International Institute for Aerospace Survey and Earth Sciences, Enschede, 243pp.

- Mérot, P., Crave, A., Cascuel-Odoux, C. and Louhala, S. (1994). Effect of saturated areas on backscattering coefficient of the ERS 1 synthetic aperture radar: First results. *Water Resour. Res.*, 30(2): 175–9.
- Merz, B. and Bárdossy, A. (1998). Effects of spatial variability on the rainfall runoff process in a small loess catchment. *J. Hydrology*, 212–213: 303–17.
- Merz, B. and Plate, E. J. (1997). An analysis of the effects of spatial variability of soil and soil moisture on runoff. *Water Resour. Res.*, 33(12): 2909–22.
- Michaud, J. D. (1992). Distributed rainfall-runoff modelling of thunderstorm generated floods: A case study in a mid-sized, semiarid watershed in Arizona. Ph.D dissertation, Department of Hydrology and Water Resources, University of Arizona, Tucson, Arizona.
- Michaud, J. D. and Shuttleworth, W. J. (1997). Aggregate descriptions of land atmosphere interactions. *J. Hydrology* (special issue), 190(3–4).
- Michaud, J. D. and Sorooshian, S. (1994a). Effect of rainfall-sampling errors on simulations of desert flash floods. *Water Resour. Res.*, 30(10): 2765–75.
- Michaud, J. D. and Sorooshian, S. (1994b). Comparison of simple versus complex distributed runoff models on a mid-sized semiarid watershed. *Water Resour. Res.*, 30(3): 593–605.
- Miljøstyrelsen (1983). Karup river investigation. An investigation of the hydrological impacts of irrigation. Miljøprojekter 51, National Agency of Environmental Protection and the counties of Viborg, Ringkjøbing and Århus (in Danish).
- Milly, P. C. D. and Eagleson, P. S. (1987). Effects of spatial variability on annual average water balance. *Water Resour. Res.*, 23(11): 2135–43.
- Milly, P. C. D. and Eagleson, P. S. (1988). Effects of storm scale on surface runoff volume. *Water Resour. Res.*, 24(4): 620–24.
- Molnar, L. (1992). *Principles for a new edition of the digital elevation modeling system, SCOP*. International Archives for Photogrammetry and Remote Sensing, XXIX/B4, Washington, pp.962–8.
- Monteith, J. L. (1995a). A reinterpretation of stomatal responses to humidity. *Plant, Cell and Environment*, 18: 357–64.
- Monteith, J. L. (1995b). Accommodation between transpiring vegetation and the convective boundary layer. *J. Hydrology*, 166: 251–63.
- Monteith, J. L. and Unsworth, M. H. (1990). *Principles of Environmental Physics*, 2nd edition. Edward Arnold, New York, 290pp.
- Moore, I. D. and Burch, G. J. (1986). Sediment transport capacity of sheet and rill flow: Application of unit stream power theory. *Water Resour. Res.*, 22: 1350–60.
- Moore, I. D., Burch, G. J. and Mackenzie, D. H. (1988a). Topographic effects on the distribution of surface soil water and the location of ephemeral gullies. *Trans. American Society of Agricultural Engineers*, 31(4): 1098–107.
- Moore, I. D. and Foster, G. R. (1990). Hydraulics and overland flow. Chapter 7 in: Anderson, M. G. and Burt, T. P. (Eds), *Process Studies in Hillslope Hydrology*. Wiley, pp.215–54.
- Moore, I. D. and Grayson, R. B. (1991). Terrain-based catchment partitioning and runoff prediction using vector elevation data. *Water Resour. Res.*, 27: 1177–91.
- Moore, I. D., Grayson, R. B. and Ladson, A. R. (1991). Digital terrain modelling: A review of hydrological, geomorphological, and biological applications. *Hydrological Processes*, 5: 3–30.
- Moore, I. D., O’Loughlin, E. M. and Burch, G. J. (1988b). A contour based topographic model for hydrological and ecological applications. *Earth Surface Processes*, 13: 305–20.
- Moore, R. D. and Thompson, J. C. (1996). Are water-table variations in a shallow forest soil consistent with the TOPMODEL concept? *Water Resour. Res.*, 32(3): 663–9.
- Moore, R. J., Harding, R. J., Austin, R. M., Bell, V. A. and Lewis, D. R. (1996). Development of improved methods for snowmelt forecasting. R&D Note 402, Research Report, Institute of Hydrology, Wallingford.
- Morton, A. (1993). Mathematical models: Questions of trustworthiness. *Brit. J. Phil. Soc.*, 44: 659–74.

- Mott, K. A. and Parkhurst, D. F. (1992). Stomatal responses to humidity in air and helox. *Plant, Cell and Environment*, 14: 509–15.
- Mroczkowski, M., Raper, G. P. and Kuczera, G. (1997). The quest for more powerful validation of conceptual catchment models. *Water Resour. Res.*, 33(10): 2325–35.
- Munley, W. G. and Hipps, L. E. (1991). Estimation of regional evaporation for a tallgrass prairie from measurements of properties of the atmospheric boundary layer. *Water Resour. Res.*, 27: 225–30.
- Myers, D. E. (1994). Spatial interpolation: An overview. *Geoderma*, 62: 17–28.
- Myrabø, S. (1986). Runoff studies in a small catchment. *Nordic Hydrology*, 17: 335–46.
- Myrabø, S. (1988). Automation in Hillslope Hydrology. NHP-rapport 22, pp.36–45.
- Myrabø, S. (1997). Temporal and spatial scale of response area and groundwater variation in till. *Hydrological Processes*, 11: 1861–80.
- Nash, J. E. and Sutcliffe, J. V. (1970). River flow forecasting through conceptual models, I. A discussion of principles. *J. Hydrology*, 10: 228–90.
- Negri, A. J., Anagnostou, E. N. and Adler, R. F. (2000). A 10-year climatology of Amazonian rainfall derived from passive microwave satellite observations. *J. of Applied Meteorology*, 39: 42–56.
- Neuman, S. P., Fogg, G. E. and Jacobson, E. A. (1980). A statistical approach to the inverse problem of aquifer hydrology: 2. Case Study. *Water Resour. Res.*, 16(1): 32–58.
- Neuman, S. P. and Yakowitz, S. (1979). A statistical approach to the inverse problem of aquifer hydrology: 1. Theory. *Water Resour. Res.*, 15(4): 845–60.
- Neuwirth, F. (1982). Beziehungen zwischen den kurzwelligen Strahlungskomponenten auf die horizontale Fläche und der Bewölkung an ausgewählten Stationen in Österreich (Relationships between the components of shortwave radiation at the horizontal surface and cloudiness at selected stations in Austria). *Archives for Meteorology, Geophysics, and Bioclimatology, Ser. B*, 30: 29–43.
- New, M., Hulme, M. and Jones, P. (1999). Representing twentieth-century space-time climate variability, Part I: Development of a 1961–90 mean monthly terrestrial climatology. *J. Climate*, 12(3): 829–56.
- Nezry, E., Lopez, A. and Touzy, R. (1991). Detection of structural and textural features for SAR images filtering. In: *Proc. International Geoscience and Remote Sensing Symposium (IGARSS)*. Espoo, Finland, pp.2169–72.
- Norman, J. M. (1993). Scaling processes between leaf and canopy levels. In: Ehleringer, J. P. and Field, C. B. (Eds), *Scaling Physiological Processes. Leaf to Globe*. Academic Press, New York, pp.41–76.
- Norman, J. M. and Becker, F. (1995). Terminology in thermal infrared remote sensing of natural surfaces. *Agric. For. Meteorol.* 77: 153–66.
- Norman, J. M., Kustas, W. P. and Humes, K. S. (1995). A two-source approach for estimating soil and vegetation energy fluxes from observations of directional radiometric surface temperature. *Agric. For. Meteorol.*, 77: 263–93.
- Northcote, K. H. (1979). *A Factual Key for the Recognition of Australian Soils*. Rellim Technical Publications, Coffs Harbour, NSW, Australia, 123pp.
- NRCS (1998). <http://www.wcc.nrcs.usda.gov/>. Snow Survey and Water Supply Forecasting Program Manager, Natural Resources Conservation Service, 101 SW Main Street, Suite 1600, Portland, Oregon 97204-3224, U.S.A.
- Obled, C. (1990). Hydrological modeling in regions of rugged relief. In: Lang, H. and Musy, A. (Eds), *Hydrology in Mountainous Regions. I – Hydrological Measurements; the Water Cycle*, Proc. Lausanne Symp., Aug. 1990, IAHS Publ. 193, 599–613.
- Obled, C. and Harder, H. (1979). A review of snowmelt in the mountain environment. In: Colbeck, S. C. and Ray, M. (Eds), *Modeling Snow Cover Runoff*. Proc. Conf. Hanover. U.S. Army CRREL, pp.179–204.
- Obled, C., Wendling, J. and Beven, K. (1994). The sensitivity of hydrologic models to spatial rainfall patterns: An evaluation using observed data. *J. Hydrology* 159: 305–33.

- O'Callaghan, J. F. and Mark, D. M. (1984). The extraction of drainage networks from digital elevation data. *Computer Vision, Graphics and Image Processing*, 28: 323–44.
- Ogden, F. L. and Julien, P. Y. (1993). Runoff sensitivity to temporal and spatial rainfall variability at runoff plane and small basin scale. *Water Resour. Res.*, 29(8): 2589–97.
- Ogden, F. L. and Julien, P. Y. (1994). Runoff model sensitivity to radar rainfall resolution. *J. Hydrology.*, 158: 1–18.
- Olyphant, G. A. (1986). Longwave radiation in mountainous areas and its influence on the energy balance of alpine snowfields. *Water Resour. Res.*, 22(1): 62–6.
- Ophori, D. U. and Tóth, J. (1989). Characterization of ground-water flow by field mapping and numerical simulation, Ross Creek Basin, Alberta, Canada. *Ground Water*, 27(2): 193–201.
- Oreskes, N., Shrader-Frechette, K. and Belitz, K. (1994). Verification, validation and confirmation of numerical models in the earth sciences. *Science*, 264: 641–6.
- Osborn, H. B. (1964). Effect of storm duration on runoff from rangeland watersheds in the semiarid southwestern United States. *Bull. IASH IX(4)*: 40–7.
- Osborn, H. B. (1977). Point to area convective rainfall simulation. *Proc. Weather–Climate Modelling for Real-Time Applications in Agriculture and Forest Meteorology*, 13th Agriculture and Forest Meteorology Conference, American Meteorological Society, pp. 51–2.
- Osborn, H. B. and Lane, L. J. (1972). Depth–area relationships for thunderstorm rainfall in Southeastern Arizona. *Trans. ASAE*, 15(4): 670–3, 680.
- Osborn, H. B. and Lane, L. J. (1981). Point–area–frequency conversions for summer rainfall in Southeastern Arizona, Hydrology and water resources in Arizona and the Southwest. Office of Arid Land Studies, Univ. of Arizona, Tucson, 11: 39–42.
- Osborn, H. B., Shirley, E. D., Davis, D. R. and Koehler, R. B. (1980). Model of time and space distribution of rainfall in Arizona and New Mexico. USDA-SEA Agricultural Reviews and Manuals, ARM-W-14, 27pp.
- Over, T. M. and Gupta, V. K. (1994). Statistical analysis of mesoscale rainfall: Dependence of a random cascade generator on large-scale forcing. *J. Appl. Meteorol.*, 33(12): 1526–42.
- Over, T. M. and Gupta, V. K. (1996). A space-time theory of mesoscale rainfall using random cascades. *J. Geophys. Res.*, 101(D21): 26319–31.
- Palacios-Vélez, O. L., Gandoy-Bernasconi, W. and Cuevas-Renaud, B. (1998). Geometric analysis of surface runoff and the computation order of unit elements in distributed hydrological models. *J. Hydrology*, 211(1–4): 266–74.
- Paniconi, C. and Wood, E. F. (1993). A detailed model for simulation of catchment scale subsurface hydrologic processes. *Water Resour. Res.*, 29(6): 1601–20.
- Parkin, G., O'Donnell, G. O., Ewen, J., Bathurst, J. C., O'Connell, P. E. and Lavabre, J. (1996). Validation of catchment models for predicting land-use and climate change impacts. 2. Case study for a Mediterranean catchment. *J. Hydrology*, 175: 595–613.
- Parsons, A. J., Abrahams, A. D. and Luk, S.-H. (1990). Hydraulics of interrill overland flow on a semi-arid hillslope, southern Arizona. *J. Hydrology*, 117: 255–73.
- Peck, A., Gorelick, S., de Marsily, G., Foster, S. and Kovalevsky, V. (1988). *Consequences of Spatial Variability in Aquifer Properties and Data Limitations for Groundwater Modelling Practice*. IAHS Publ. No. 175, IAHS Press, Wallingford.
- Peck, E. L. and Schaake, J. C. (1990). Network design for water supply forecasting in the West. *Water Resour. Bull.*, 26(1): 87–99.
- Perica, S. (1995). A model for multiscale disaggregation of spatial rainfall based on coupling meteorological and scaling descriptions. PhD Thesis, University of Minnesota, Minneapolis.
- Perica, S. and Foufoula-Georgiou, E. (1996a). Linkage of scaling and thermodynamic parameters of rainfall: Results from midlatitude mesoscale convective systems. *J. Geophys. Res.*, 101(D3): 7431–48.

- Perica, S. and Foufoula-Georgiou, E. (1996b). A model for multiscale disaggregation of spatial rainfall based on coupling meteorological and scaling descriptions. *J. Geophys. Res.*, 101(D21): 26347–61.
- Pessoa, M. L., Bras, R. L. and Williams, E. R. (1993). Use of weather radar for flood forecasting in the Sieve River Basin: A sensitivity analysis. *J. Appl. Meteorol.*, 32(3): 462–75.
- Philip, J. R. (1957). The theory of infiltration. The infiltration equation and its solution. *Soil Science*, 83: 345–57.
- Philip, J. R. (1975). Some remarks on science in catchment prediction. In: Chapman, T. G. and Dunin, F. X. (Eds), *Prediction in Catchment Hydrology: A National Symposium on Hydrology*. Australian Academy of Science, pp.23–30.
- Pilgrim, D. H. and Cordery, I. (1975). Rainfall temporal patterns for design floods. *J. Hydr. Div., ASCE*, 101(HY1): 81–95.
- Pitman, A. J., Henderson-Sellers, A. and Yang, Z. L. (1990). Sensitivity of regional climates to localized precipitation in global models. *Nature*, 346: 734–7.
- Pomeroy, J. W. and Gray, D. M. (1995). Snowcover: accumulation, relocation and management. National Hydrology Research Institute Science Report No. 7, Saskatoon, Saskatchewan, Canada, 144pp.
- Popper, K. (1977). *The Logic of Scientific Discovery*. Routledge, 480pp.
- Porcello, L., Massey, N., Innes, R. and Marks, J. (1976). Speckle reduction in synthetic aperture radars. *J. Opt. Soc. Am.*, 66(11): 1305–11.
- Press, W. H., Teukolsky, S. A., Vetterling, W. T. and Flannery, B. P. (1986). *Numerical Recipes in FORTRAN: The Art of Scientific Computing*, 2nd Edition. Cambridge University Press.
- Price, A. G. and Dunne, T. (1976). Energy balance computations of snowmelt in a subarctic area. *Water Resour. Res.*, 12(4): 686–94.
- Price, J. C. (1990). Using spatial context in satellite data to infer regional scale evapotranspiration. *IEEE Trans. Geosci. Remote Sens.*, GE-28: 940–8.
- Priestley, C. H. B. and Taylor, R. J. (1972). On the assessment of the surface heat flux and evaporation using large-scale parameters. *Mon. Weather Rev.*, 100: 81–92.
- Puckett, W. E., Dane, J. H. and Hajek, B. F. (1985). Physical and mineralogical data to determine soil hydraulic properties. *Soil Sci. Soc. Am. J.*, 49: 831–6.
- Quinn, P. and Anthony, S. (in review). Digital terrain analysis and hydrological modelling. In: *Spatial Analysis and Hydrological Modelling*. Wiley.
- Quinn, P. F. and Beven, K. J. (1993). Spatial and temporal predictions of soil moisture dynamics, runoff, variable source areas and evapotranspiration for Plynlimon, Mid-Wales. *Hydrological Processes*, 7: 425–48.
- Quinn, P. F., Beven, K., Chevallier, P. and Planchon, O. (1991). The prediction of hillslope flow paths for distributed hydrological modelling using digital terrain models. *Hydrological Processes*, 5: 59–79.
- Quinn, P. F., Beven, K. J. and Lamb, R. (1995). The $\ln(a/\tan \beta)$ index: how to calculate it and how to use it within the TOPMODEL framework. *Hydrological Processes*, 9(2): 161–82.
- Rawls, W. J., Brakensiek, D. L. and Miller, N. (1983). Green-Ampt infiltration parameters from soils data. *J. Hydraul. Eng.*, 109: 62–70.
- Refsgaard, J. C. (1996). Terminology, modelling protocol and classification of hydrological model codes. In: Refsgaard, J. C. and Abbott, M. B. (Eds), *Distributed Hydrological Modelling*, Kluwer, Dordrecht, The Netherlands, pp.17–40.
- Refsgaard, J. C. (1997). Parameterisation, calibration and validation of distributed hydrological models. *J. Hydrology*, 198(1–4): 69–97.
- Refsgaard, J. C. and Knudsen, J. (1996). Operational validation and intercomparison of different types of hydrological models. *Water Resour. Res.*, 32(7): 2189–202.
- Refsgaard, J. C. and Storm, B. (1995). MIKE SHE. In: Singh, V. P. (Ed.), *Computer Models of Watershed Hydrology*, Water Resources Publications, Highlands Ranch, Colorado, pp.809–46.

- Refsgaard, J. C. and Storm, B. (1996). Construction, calibration and validation of hydrological models. In: Abbott, M. B. and Refsgaard, J. C. (Eds), *Distributed Hydrological Modelling*. Kluwer, Dordrecht, The Netherlands, pp.41–54.
- Refsgaard, J. C., Storm, B. and Abbott, M. B. (1996). Comment on: “A discussion of distributed hydrological modelling” by K. Beven. In: Refsgaard, J. C. and Abbott, M. B. (Eds), *Distributed Hydrological Modelling*. Kluwer, Dordrecht, The Netherlands, pp.279–87.
- Renard, K. G., Lane, L. J., Simanton, J. R., Emmerich, W. E., Stone, J. J., Weltz, M. A., Goodrich, D. C. and Yakowitz, D. S. (1993). Agricultural impacts in an arid environment: Walnut Gulch case study. *Hydrol. Sci. Tech.*, 9(1–4): 145–90.
- Rhea, J. O. (1978). Orographic precipitation model for hydrometeorological use. Ph.D Thesis, Colorado State University, Fort Collins, Colorado.
- Richards, J. A. (1986). *Remote Sensing Digital Image Analysis*. Springer-Verlag.
- Rieger, W. (1998). A phenomenon-based approach to upslope contributing area and depressions in DEMs. *Hydrological Processes*, 12: 857–72.
- Rignot, E. J. M. and van Zyl, J. J. (1993). Change detection techniques for ERS-1 SAR data. *IEEE Trans. Geosc. Rem. Sens.*, 31(4): 896–906.
- Riley, S. J., Crozier, P. and Blong, R. J. (1981). An inexpensive and easily installed runoff plot. *J. Soil Conservation Service, New South Wales*, 37: 144–7.
- Robins, J. S., Kelly, L. L. and Hamon, W. R. (1965). Reynolds Creek in Southwest Idaho: An outdoor hydrologic laboratory. *Water Resour. Res.*, 1(3): 407–13.
- Rodhe, A. (1981). Spring flood meltwater or ground-water? *Nordic Hydrology*, 12(1): 21–30.
- Rohrer, M. B. (1992). Die Schneedecke im Schweizer Alpenraum und ihre Modellierung. (Snow cover modelling in the Swiss Alps.) Zürcher Geographische Schriften, Heft 49. Geographisches Institut der Eidgenössischen Technischen Hochschule Zürich, 178pp.
- Romano, N. and Santini, A. (1997). Effectiveness of using pedo-transfer functions to quantify the spatial variability of soil water retention characteristics. *J. Hydrology*, 202: 137–57.
- Romanowicz, R., Beven, K. J. and Tawn, J. A. (1994). Evaluation of predictive uncertainty in nonlinear hydrological models using a Bayesian approach. In: Barnett, V. and Feridum Turkman, K. (Eds), *Statistics for the Environment 2: Water Related Issues*, pp.297–317.
- Rosenzweig, C. and Abramopoulos, F. (1997). Land-surface model development for the GISS GCM. *J. Climate*, 10(8): 2040–54.
- Rosso, R. (1994). An introduction to spatially distributed modelling of basin response. In: Rosso, R., Peano, A., Becchi, I. and Bemporad, G. A. (Eds), *Advances in Distributed Hydrology*. Water Resources Publications, Highlands Ranch, Colorado, pp.3–30.
- Roth, M. and Oke, T. R. (1995). Relative efficiencies of turbulent transfer of heat, mass, and momentum over a patchy urban surface. *J. Atmos. Sci.*, 52: 1863–74.
- Salahshour Dehchali, J. (1993). Etude hydrologique du bassin versant du Coët-dan à Naizin; cartographie et modélisation des zones humides. Mémoire présenté en vue de l’obtention du Diplôme d’Agronomie Approfondie, INRA-ENSAR, 43pp.
- Salvucci, G. D. (1998). Limiting relations between soil moisture and texture with implications for measured, modelled and remotely sensed estimates. *Geophys. Res. Letters*, 25(10): 1757–60.
- Salvucci, G. D. and Entekhabi, D. (1994a). Equivalent steady moisture profile and the time compression approximation in water balance modelling. *Water Resour. Res.*, 30(10): 2737–49.
- Salvucci, G. D. and Entekhabi, D. (1994b). Comparison of the Eagleson statistical-dynamical water balance model with numerical simulations. *Water Resour. Res.*, 30(10): 2751–7.
- Salvucci, G. D. and Entekhabi, D. (1995). Hillslope and climatic controls on hydrologic fluxes. *Water Resour. Res.*, 31(7): 1725–39.
- Salvucci, G. D. and Entekhabi, D. (1997). Corrections to “Hillslope and climatic controls on hydrologic fluxes”. *Water Resour. Res.*, 33(1): 277.
- Satterlund, D. R. (1979). An improved equation for estimating long-wave radiation from the atmosphere. *Water Resour. Res.*, 15: 1643–50.

- Saulnier, G.-M., Beven, K. and Obled, C. (1997c). Digital elevation analysis for distributed hydrological modelling: Reducing scale dependence in effective hydraulic conductivity values. *Water Resour. Res.*, 33(9): 2097–101.
- Saulnier, G.-M., Beven, K. and Obled, C. (1997a). Including spatially variable effective soil depths in TOPMODEL. *J. Hydrology*, 202(1–4): 158–72.
- Saulnier, G. M., Obled, C. and Beven, K. (1997b). Analytical compensation between DTM grid resolution and effective values of saturated hydraulic conductivity within the TOPMODEL framework. *Hydrological Processes*, 11(9): 1331–46.
- Schertzer, D. and Lovejoy, S. (1987). Physical modeling and analysis of rain and clouds by anisotropic scaling multiplicative processes, *J. Geophys. Res.*, 92(D8): 9693–714.
- Schlesinger, S., Crosbie, R. E., Gagné, R. E., Innis, G. S., Lalwani, C. S., Loch, J., Sylvester, J., Wright, R. D., Kheir, N. and Bartos, D. (1979). Terminology for model credibility. SCS Technical Committee on Model Credibility. *Simulation*, 32(3): 103–4.
- Schmid, H. P. (1994). Source areas for scalars and scalar fluxes. *Boundary-Layer Meteorol.*, 67: 293–318.
- Schmugge, T., Jackson, T. J., Kustas, W. P., Roberts, R., Parry, R., Goodrich, D. C., Amer, S. A. and Wertz, M. A. (1994). Push broom microwave observations of surface soil moisture in Monsoon '90. *Water Resour. Res.*, 30(5): 1321–7.
- Schmugge, T. J., Kustas, W. P. and Humes, K. S. (1998). Monitoring land surface fluxes using ASTER observations. *IEEE Trans. Geosci. Remote Sens.*, 36(5): 1–10.
- Schuepp, P. H., Leclerc, M. Y., Macpherson, J. I. and Desjardins, R. L. (1990). Footprint prediction of scalar fluxes from analytical solutions of the diffusion equation. *Boundary-Layer Meteorol.*, 50: 355–73.
- Schulz, G. A. (1988). Remote sensing in hydrology. *J. Hydrology*, 100: 239–65.
- Seaman, N. L. (1990). Newtonian Nudging: A Four-Dimensional Approach to Data Assimilation. Mesoscale Data Assimilation, 1990 Summer Colloquium, National Center for Atmospheric Research, Boulder, Colorado, 6 June–3 July.
- Seed, A. W., Srikanthan, R. and Menabde, M. (1999). A space and time model for design storm rainfall. *J. Geophys. Res.* 104 (D24): 31623–30.
- Seibert, J., Bishop, K. H. and Nyberg, L. (1997). A test of TOPMODEL's ability to predict spatially distributed groundwater levels. *Hydrological Processes*, 11(9): 1131–44.
- Sellers, P. J., Heiser, M. D., Hall, F. G., Goetz, S. J., Strebel, D. E., Verma, S. B., Desjardins, R. L., Schuepp, P. H. and MacPherson, J. I. (1995). Effects of spatial variability in topography, vegetation cover and soil moisture on area-averaged surface fluxes: A case study using FIFE 1989 data. *J. Geophys. Res.*, 100(D12): 25607–29.
- Seyfried, M. S. and Wilcox, B. P. (1995). Scale and the nature of spatial variability: Field examples having implications for hydrologic modelling. *Water Resour. Res.*, 31(1): 173–84.
- Shaw, E. M. and Lynn, P. P. (1972). Areal rainfall evaluation using two surface fitting techniques. *Bull. Int. Assoc. Hydrol. Sci.*, XVII(4) 12: 419–33.
- Shen, H. W. and Julien, P. Y. (1992). Erosion and sediment transport. Chapter 12 in: Maidment, D. (Ed.), *Handbook of Hydrology*. McGraw-Hill, New York, pp.12.1–12.61.
- Shi, Z. and Fung, K. B. (1994). A comparison of digital speckle filters. In: *Proc. International Geoscience and Remote Sensing Symposium (IGARSS)*. Pasadena, USA, pp.2129–33.
- Shuttleworth, W. J. (1991). Insight from large-scale observational studies of land/atmosphere interactions. *Surv. Geophys.*, 29: 585–606.
- Shuttleworth, W. J. and Wallace, J. S. (1985). Evaporation from sparse crops – An energy combination theory. *Quart. J. R. Meteorol. Soc.*, 111: 839–55.
- Singh, A. (1989). Digital change detection techniques using remotely-sensed data. *Int. J. Rem. Sens.*, 10(6): 989–1003.
- Singh, V. P. (Ed.) (1995). *Computer Models of Watershed Hydrology*. Water Resources Publications, Highlands Ranch, Colorado, 1130pp.

- Sivapalan, M., Beven, K. J. and Wood, E. F. (1987). On hydrologic similarity, 2, A scaled model of storm runoff production. *Water Resour. Res.*, 23(2): 2266–78.
- Sivapalan, M. and Blöschl, G. (1998). Transformation of point rainfall to areal rainfall: intensity–duration–frequency curves. *J. Hydrology*, 204: 150–67.
- Sivapalan, M. and Woods, R. A. (1995). Evaluation of the effects of general circulation model's subgrid variability and patchiness of rainfall and soil moisture on land surface water balance fluxes. In: Kalma, J. D. and Sivapalan, M. (Eds), *Scale issues in Hydrological Modelling*. John Wiley, New York, pp.453–73.
- Smith, M., Allen, R., Monteith, J. L., Perrier, A., Santos Pereira, L. and Segeren, A. (1992). Expert consultation on revision of FAO methodologies for crop water requirements. Land and Water Development Division, Food and Agriculture Organization of the United Nations, Rome.
- Smith, R. E., Goodrich, D. C. and Woolhiser, D. A. (1990). Areal effective infiltration dynamics for runoff on small catchments. *Trans. 14th Inter. Congress of Soil Sci.*, Volume I: Commission I, Kyoto, Japan, Aug. 1990, pp.22–7.
- Smith, R. E., Goodrich, D. R., Woolhiser, D. A. and Simanton, J. R. (1994). Comment on “Physically based hydrological modelling 2: Is the concept realistic?” by Grayson, Moore and McMahon. *Water Resour. Res.*, 30: 851–4.
- Smith, R. E., Goodrich, D. R., Woolhiser, D. A. and Unkrich, C. L. (1995). KINEROS – A KINematic Runoff and EROSION Model. Chapter 20 in: Singh, V. P. (Ed.), *Computer Models of Watershed Hydrology*. Water Resources Publications, Highlands Ranch, Colorado, pp.697–732.
- Smith, R. E. and Hebbert, R. H. B. (1979). A Monte Carlo analysis of the hydrologic effects of spatial variability of infiltration. *Water Resour. Res.*, 15(2): 419–29.
- Smith, R. E. and Parlange, J.-Y. (1978). A parameter-efficient hydrologic infiltration model. *Water Resour. Res.*, 14(33): 533–8.
- Sorooshian, S. and Dracup, J. (1980). Stochastic parameter estimation procedures for hydrological rainfall–runoff models: correlated and heteroscedastic error cases. *Water Resour. Res.*, 16(2): 430–42.
- Sorooshian, S. and Gupta, V. K. (1995). Model calibration. Chapter 2 in: Singh, V. P. (Ed.), *Computer Models of Watershed Hydrology*. Water Resources Publications, Highlands Ranch, Colorado, pp.23–63.
- Sorooshian, S., Gupta, H. V. and Rodda, J. C. (1997). *Land Surface Processes in Hydrology – Trials and Tribulations of Modeling and Measuring*. Springer, Berlin, 497pp.
- Stannard, D. I. (1993). Comparison of Penman–Monteith, Shuttleworth–Wallace, and modified Priestley–Taylor evapotranspiration models for wildland vegetation in semiarid rangeland. *Water Resour. Res.*, 29: 1379–92.
- Stannard, D. I., Blanford, J. H., Kustas, W. P., Nichols, W. D., Amer, S. A., Schmutge, T. J. and Weltz, M. A. (1994). Interpretation of surface flux measurements in heterogeneous terrain during Monsoon '90. *Water Resour. Res.*, 30(5): 1227–39.
- Stauffer, D. R. and Seaman, N. L. (1990). Use of four-dimensional data assimilation in a limited-area mesoscale model. Part I: Experiments with synoptic-scale data. *Mon. Weather Rev.*, 118: 1250–77.
- Stendal, M. M. (1978). Hydrological data – Norden, Karup representative basin, Denmark. Data volume 1965–77. Danish National Committee for the IHD, Copenhagen.
- Stephenson, G. R. and Freeze, R. A. (1974). Mathematical simulation of subsurface flow contributions to snowmelt and runoff, Reynolds Ck. Watershed, Idaho. *Water Resour. Res.*, 10: 284–94.
- Stewart, J. B. and Verma, S. B. (1992). Comparison of surface fluxes and conductances at two contrasting sites within the FIFE area. *J. Geophys. Res.*, 97: 18623–8.
- Stoertz, M. W. and Bradbury, K. R. (1989). Mapping recharge areas using a ground-water flow model – A case study. *Ground Water*, 27(2): 220–8.

- Stone, J. J., Lane, L. J., Shirley, E. D. and Renard, K. G. (1986). A runoff-sediment yield model for semiarid regions. *Proc. 4th Fed. Interagency Sedimentation Conf.*, Las Vegas, NV, pp. 6-75-6-84.
- Stull, R. (1988). *An Introduction to Boundary Layer Meteorology*. Kluwer, Boston, MA.
- Styczen, M. (1995). Validation of pesticide leaching models. In: *Leaching models and EU registration*. Final report of the work of the Regulatory Modelling Work Group of FOCUS (FORum for the Co-ordination of pesticide fate models and their Use), DOC.4592/VI/95, European Commission.
- Styczen, M. and Storm, B. (1993). Modelling of N-movements on catchment scale – a tool for analysis and decision making. 1. Model description & 2. A case study. *Fertilizer Research*, 36: 1–17.
- Sugita, M. and Brutsaert, W. (1991). Daily evaporation over a region from lower boundary layer profiles. *Water Resour. Res.*, 27: 747–52.
- Swiatek, E. (1992). Estimating regional surface fluxes from measured properties of the atmospheric boundary layer in a semiarid ecosystem. M.S. Thesis, Utah State University, Logan, UT.
- Syed, K. H. (1994). Spatial Storm Characteristics and Basin Response. M.S. Thesis, Department of Hydrology and Water Resources, The University of Arizona, Tucson, Arizona, 261pp.
- Tabios, G. Q. and Salas, J. D. (1985). A comparative analysis of techniques for spatial interpolation of precipitation. *Water Resour. Bull.*, 21: 365–80.
- Talsma, T. (1987). Re-evaluation of the well permeameter as a method for measuring hydraulic conductivity. *Australian J. Soil Research*, 25: 361–8.
- Talsma, T. and Hallam, P. M. (1980). Hydraulic conductivity measurements of forest catchments. *Australian J. Soil Research*, 18: 139–48.
- Tanner, M. H. (1992). *Tools for statistical inference: Observed data and data augmentation methods*. Lecture notes in statistics, 67, Springer-Verlag.
- Tarboton, D. G. (1997). A new method for the determination of flow directions and upslope areas in grid digital elevation models. *Water Resour. Res.*, 33(2): 309–19.
- Tarboton, D. G., Chowdhury, T. G. and Jackson, T. H. (1995). A spatially distributed energy balance snowmelt model. In: Tonnessen, K. A. et al. (Eds), *Biogeochemistry of Seasonally Snow-Covered Catchments*, Proc. Boulder Symposium, July 3–14, IAHS Publ. no. 228, pp.141–55.
- Tarboton, D. G. and Luce, C. H. (1996). Utah Energy Balance Snow Accumulation and Melt Model (UEB), Computer model technical description and user's guide, Utah Water Research Laboratory and USDA Forest Service Intermountain Research Station (<http://www.engineering.usu.edu/dtarb/>).
- Tessier, Y., Lovejoy, S. and Schertzer, D. (1993). Universal multifractals: Theory and observations for rain and clouds. *J. Appl. Meteor.*, 32(2): 223–50.
- Thiessen, A. H. (1911). Precipitation for large areas. *Monthly Weather Review*, 39: 1082–4.
- Tóth, J. (1962). A theory of groundwater motion in small drainage basins in central Alberta, Canada. *J. Geophys. Res.*, 67(11): 4375–87.
- Tóth, J. (1963). A theoretical analysis of groundwater flow in small drainage basins. *J. Geophys. Res.*, 68(16): 4795–812.
- Tóth, J. (1966). Mapping and interpretation of field phenomena for groundwater reconnaissance in a prairie environment, Alberta, Canada. *Bull. IASH*, 11(2): 20–68.
- Tsang, C.-F. (1991). The modeling process and model validation. *Ground Water*, 29: 825–31.
- Turner, J. V. and Macpherson, D. K. (1987). Mechanisms affecting streamflow and stream water quality: an approach via stable isotope, hydrogeochemical, and time series analysis. *Water Resour. Res.*, 26(12): 3005–19.
- Tyndale-Biscoe, J. P., Moore, G. A. and Western, A. W. (1998). A system for collecting spatially variable terrain data. *Computers and Electronics in Agriculture*, 19(2): 113–28.

- U.S. Army Corps of Engineers (1956). Snow Hydrology, Summary report of the Snow Investigations, U.S. Army Corps of Engineers, North Pacific Division, Portland, Oregon.
- U.S. Senate Committee (1959). Report on findings on facility needs for soil and water research. 175pp.
- Ulaby, F. T., Dubois, P. C. and van Zyl, J. (1996). Radar mapping of surface soil moisture. *J. Hydrology*, 184: 57–84.
- Ulaby, F. T., Moore, R. K. and Fung, A. K. (1982). *Microwave Remote Sensing, Active and Passive, Vol. II: Radar Remote Sensing and Surface Scattering and Emission Theory*, Artech House, Inc.
- van Straten, G. and Keesman, K. J. (1991). Uncertainty propagation and speculation in projective forecasts of environmental change: A lake-eutrophication example. *J. Forecasting*, 10: 163–90.
- Valdes, J. B., Rodriguez-Iturbe, I. and Gupta, V. K. (1985). Approximations of temporal rainfall from a multidimensional model. *Water Resour. Res.*, 21(8): 1259–70.
- Vanmarcke, E. (1983). *Random Fields: Analysis and Synthesis*. MIT Press, Cambridge, Massachusetts, 382pp.
- Venugopal, V. (1999). Spatio-temporal organization and space-time downscaling of precipitation fields. PhD Thesis, University of Minnesota, Minneapolis.
- Venugopal, V., Foufoula-Georgiou, E. and Sapozhnikov, V. (1999a). Evidence of dynamic scaling in space-time rainfall. *J. Geophys. Res.*, 104(D24): 31599–610.
- Venugopal, V., Foufoula-Georgiou, E. and Sapozhnikov, V. (1999b). A space-time downscaling model for rainfall. *J. Geophys. Res.*, 104(D16): 705–21.
- Verhoef, A., de Bruin, H. A. R. and Van Den Hurk, B. J. J. M. (1997). Some practical notes on the parameter kB-1 for sparse vegetation. *J. Appl. Meteorol.*, 36: 560–72.
- Verhoest, N. E. C., Troch, P. A., Paniconi, C. and De Troch, F. P. (1998). Mapping basin scale variable source areas from multitemporal remotely sensed observations of soil moisture behavior. *Water Resour. Res.*, 34(12): 3235–44.
- Verstraete, M. M. (1988). Radiation transfer in plant canopies: Scattering of solar radiation and canopy reflectance. *J. Geophys. Res.*, 93(D8): 9483–94.
- Verstraete, M. M. (1987). Radiation transfer in plant canopies: Transmission of direct solar radiation and the role of leaf orientation. *J. Geophys. Res.*, 92(D9): 10985–95.
- Verstraete, M. M., Pinty, B. and Dickinson, R. E. (1990). A physical model of the bidirectional reflectance of vegetation canopies, 1. Theory. *J. Geophys. Res.*, 95(D8): 11755–65.
- Vertessy, R. A. and Elsenbeer, H. (1999). Distributed modeling of storm flow generation in an Amazonian rain forest catchment: Effects of model parameterization. *Water Resour. Res.*, 35(7): 2173–87.
- Vertessy, R. A., Hatton, T. J., Benyon, R. J. and Dawes, W. R. (1996). Long term growth and water balance predictions for a mountain ash (*E. regnans*) forest subject to clearfelling and regeneration. *Tree Physiol.*, 16: 221–32.
- Vertessy, R. A., Hatton, T. J., O'Shaughnessy, P. J. and Jayasuriya, M. D. A. (1993). Predicting water yield from a mountain ash forest catchment using a terrain analysis-based catchment model. *J. Hydrology*, 150: 665–700.
- Vickers, D. and Mahrt, L. (1997). Quality control and flux sampling problems for tower and aircraft data. *J. Atmos. Oceanic Tech.*, 14: 512–26.
- Viessman, Jr., W. and Lewis, G. L. (1996). *Introduction to Hydrology*. Harper Collins, New York.
- Wagner, W. (1998). Soil moisture retrieval from ERS scatterometer data. Research Report EUR 18670, European Commission Joint Research Centre, Ispra, Italy, 101pp.
- Wahba, G. and Wendelberger, J. (1980). Some new mathematical methods for variational objective analysis using splines and cross validation. *Monthly Weather Review*, 108: 36–57.
- Wang, Q. J., McConachy, F. L. N., Chiew, F. H. S., James, R., de Hoedt, G. C. and Wright, W. J. (1998). *Climatic Atlas of Australia: Maps of Evapotranspiration*. To be published by the Bureau of Meteorology, Australia.

- Wang, W. and Seaman, N. L. (1997). A comparison study of convective parameterization schemes in a mesoscale model. *Mon. Weath. Rev.*, 125: 252–78.
- Warner, T. T. and Hsu, H.-M. (2000). Nested model simulation of moist convection: the impact of coarse-grid parameterized convection on fine-grid resolved convection. *Mon. Weath. Rev.*, 128(7): 2211–31.
- Warrick, A. W., Zhang, R., Moody, M. M. and Myers, D. E. (1990). Kriging versus alternative interpolators: errors and sensitivity to model inputs. In: Roth, K. et al. (Eds), *Field-scale Water and Solute Flux in Soils*. Birkhäuser Verlag, Basel, pp.157–64.
- Watson, F. G. R., Grayson, R. B., Vertessy R. A. and McMahon, T. A. (1998). Large scale distribution modelling and the utility of detailed data. *Hydrological Processes*, 12(6): 873–88.
- Waymire, E., Gupta, V. K. and Rodriguez-Iturbe, I. (1984). A spectral theory of rainfall intensity at the meso- β scale. *Water Resour. Res.*, 20: 1453–65.
- Weisman, M. L. and Klemp, J. B. (1982). The dependence of numerically simulated convective storms on vertical wind shear and buoyancy. *Mon. Wea. Rev.*, 110: 504–20.
- Wen, X.-H. and Gómez-Hernández, J. J. (1996). Upscaling hydraulic conductivities in heterogeneous media: An overview. *J. Hydrology*, 183: ix–xxxii.
- Western, A. W. and Blöschl, G. (1999). On the spatial scaling of soil moisture. *J. Hydrology*, 217: 203–24.
- Western, A. W., Blöschl, G. and Grayson, R. B. (1998a). Geostatistical characterisation of soil moisture patterns in the Tarrawarra Catchment. *J. Hydrology*, 205: 20–37.
- Western, A. W., Blöschl, G. and Grayson, R. B. (1998b). How well do indicator variograms capture the spatial connectivity of soil moisture? *Hydrological Processes*, 12: 1851–68.
- Western, A. W., Blöschl, G. and Grayson, R. B. (2000). Towards capturing hydrologically significant connectivity in spatial patterns. *Water Resour. Res.*, in press.
- Western, A. W. and Grayson, R. B. (1998). The Tarrawarra data set: Soil moisture patterns, soil characteristics and hydrological flux measurements. *Water Resour. Res.*, 34(10): 2765–8.
- Western, A. W., Grayson, R. B., Blöschl, G., Willgoose, G. R. and McMahon, T. A. (1999a). Observed spatial organisation of soil moisture and its relation to terrain indices. *Water Resour. Res.*, 35(3): 797–810.
- Western, A. W., Grayson, R. B. and Green, T. R. (1999b). The Tarrawarra Project: High resolution spatial measurement, modelling and analysis of soil moisture and hydrological response. *Hydrological Processes*, 13(5):689–700.
- Whelan, M. J. and Anderson, J. M. (1996). Modelling spatial patterns of throughfall and interception loss in a Norway spruce (*Picea abies*) plantation at the plot scale. *J. Hydrology*, 186: 335–54.
- Wiener, N. (1949). *Extrapolation, Interpolation and Smoothing of Stationary Time Series*. MIT Press, Cambridge, Massachusetts.
- Wigmosta, M. S. and Lettenmaier, D. P. (1999). A comparison of simplified methods for routing topographically driven subsurface flow. *Water Resour. Res.*, 35(1): 255–64.
- Wigmosta, M. S., Vail, L. W. and Lettenmaier, D. P. (1994). A distributed hydrology–vegetation model for complex terrain. *Wat. Resour. Res.*, 30(6): 1665–79.
- Wilks, D. S. (1995). *Statistical Methods in the Atmospheric Sciences: An Introduction*. Academic Press, 464 pp.
- Willgoose, G. and Kuczera, G. (1995). Estimation of sub-grid scale kinematic wave parameters for hillslopes. *Hydrological Processes*, 9: 469–82.
- Williams, A. G. and Hacker, J. M. (1993). Interactions between coherent eddies in the lower convective boundary layer. *Boundary-Layer Meteorol.*, 64: 55–74.
- Williams, K. S. and Tarboton, D. G. (1999). The ABC's of snowmelt: A topographically factorized energy component snowmelt model. *Hydrological Processes*, 13: 1905–20.
- Williams, R. E. (1988). Comment on “Statistical theory of groundwater flow and transport: pore to laboratory, laboratory to formation, and formation to regional scale” by Gedeon Dagan. *Water Resour. Res.*, 24: 1197–200.

- Winchell, M., Gupta, H. V. and Sorooshian, S. (1998). On the simulation of infiltration- and saturation-excess runoff using radar-based rainfall estimates: effects of algorithm uncertainty and pixel aggregation. *Water Resour. Res.*, 34(10): 2655–70.
- Winstral, A., Elder, K. and Davis, R. (1999). Implementation of digital terrain analysis to capture the effects of wind redistribution in spatial snow modelling. In: 67th Western Snow Conference, South Lake Tahoe, CA.
- Wiscombe, W. J. and Warren, S. G. (1981). A model of the spectral albedo of snow. I: Pure snow. *J. Atmos. Sci.*, 37: 2712–33.
- WMO (1975). Intercomparison of conceptual models used in operational hydrological forecasting. WMO operational hydrology report no 7, WMO no 429. World Meteorological Organisation, Geneva.
- WMO (1986). Intercomparison of models for snowmelt runoff. WMO operational hydrology report no 23, WMO no 646. World Meteorological Organisation, Geneva.
- WMO (1992). Simulated real-time intercomparison of hydrological models. WMO operational hydrology report no 38, WMO no 779. World Meteorological Organisation, Geneva.
- Wolock, D. M. and Price, C. V. (1994). Effects of digital elevation model map scale and data resolution on a topography-based watershed model. *Water Resour. Res.*, 30: 3041–52.
- Woo, M., Heron, R., Marsh, P. and Steer, P. (1983). Comparison of weather station snowfall with winter snow accumulation in high arctic basins. *Atmos. Ocean*, 21(3): 312–25.
- Wood, E. F., Lettenmaier, D. P. and Zartarian, V. G. (1992). A land-surface hydrology parameterization with subgrid variability for general circulation models. *J. Geophys. Res.*, 97(D3): 2717–28.
- Wood, E. F., Sivapalan, M., Beven, K. and Band, L. (1988). Effects of spatial variability and scale with implications to hydrologic modeling. *J. Hydrology*, 102: 29–47.
- Woods, R., Sivapalan, M. and Duncan, M. (1995). Investigating the Representative Elementary Area concept: An approach based on field data. *Hydrological Processes*, 9: 291–312.
- Woods, R. A., Sivapalan, M. and Robinson, J. (1997). Modelling the spatial variability of subsurface runoff using a topographic index. *Water Resour. Res.*, 33(5): 1061–73.
- Woolhiser, D. A. and Goodrich, D. C. (1988). Effect of storm rainfall intensity patterns on surface runoff. *J. Hydrology*, 102: 335–54.
- Xue, M., Droegemeier, K. K., Wong, V., Shapiro, A. and Brewster, K. (1995). ARPS Version 4.0 User's Guide, 380pp. Available from the Center for Analysis and Prediction of Storms, 100 East Boyd Street, Norman, OK 73019.
- Xue, M., Droegemeier, K. K. and Wong, V. (2000a). The Advanced Regional Prediction System (ARPS) – A multiscale nonhydrostatic atmospheric simulation and prediction tool. Part I: Model dynamics and verification. *Meteor. and Atmos. Physics*, in press.
- Xue, M., Droegemeier, K. K., Wong, V., Shapiro, A., Brewster, K., Carr, F., Weber, D., Liu, Y. and Wang, D.-H. (2000b). The Advanced Regional Prediction System (ARPS) – A multiscale nonhydrostatic atmospheric simulation and prediction tool. Part II: Model physics and applications. *Meteor. and Atmos. Physics*, in press.
- Yang, D. and Woo, M. (1999). Representativeness of local snow data for large scale hydrologic investigations. *Hydrological Processes*, 13(12–13): 1977–88.
- Zak, S. K., Beven, K. J. and Reynolds, B. (1997). Uncertainty in the estimation of critical loads: a practical methodology. *Water, Air and Soil Pollution*, 98: 297–316.
- Zepeda-Arce, J., Foufoula-Georgiou, E., Droegemeier, K. (2000). Space-time rainfall organization and its role in validating quantitative precipitation forecast. *J. Geophys. Res.* 105(D8), 10129–46.
- Zermeño-Gonzalez, A. and Hipps, L. E. (1997). Downwind evolution of surface fluxes over a vegetated surface during local advection of heat and saturation deficit. *J. Hydrology*, 192: 189–210.
- Zhang, S. and Foufoula-Georgiou, E. (1997). Subgrid scale rainfall variability and effects on atmospheric and surface variable prediction. *J. Geophys. Res.*, 102(D16): 19559–73.

- Zhang, W. and Montgomery, D. R. (1994). Digital elevation model grid size, landscape representation, and hydrologic simulations. *Water Resour. Res.*, 30: 1019–28.
- Zhu, H. and Journel, A. G. (1993). Formatting and integrating soft data: stochastic imaging via the Markov–Bayes algorithm. In: Soares, A. (Ed.), *Geostatistics Tróia '92*. Kluwer, Dordrecht, The Netherlands, pp.1–12.

Index

Page numbers referring to tables are in *italics*; those referring to figures are in ***bold italics***.

- Aerial photographs 165, 169, **174**
- Aerodynamic resistance 107
- Aggregation 24, 37–38, 85, **86**, 115–117, 121–122, 155, 347
- Albedo 161, 173, 175
- Alpine catchments 159, 168
- Antecedent conditions
 - discharge 260, 261
 - wetness 76, 138, 151, 155, **155**, 187, 209, 248
- Areal reduction (of precipitation) 82, 140
- Assimilation – *see* Data assimilation
- Atmospheric boundary layer (ABL) 108–109, 111
- Auxiliary data – *see* Surrogate data
- Averaging – *see* Aggregation

- Backscattering coefficient **194**, 196, **205**, **207**
- Bedrock 9, 274, 306, 314
- Bias 23, 77, 184, 286
- Black box/empirical models 54, 164–165
- Boreholes – *see* Piezometers
- Bowen ratio 109–110
- Brightness temperature 115, 118, 135, **140**

- Calibration
 - in remote sensing 24, 136–138, **140**, 195–197, 332, 341
 - of hydrologic models 75–78, 152–153, 283–286, 288–291, 336, 338, 346
- CAPE (Convective Available Potential Energy) 87, **88**, 92, 102
- Capillary rise 310, 312
- Channel flow 68, **68**, 151–152
- Channel transmission loss 138–139, **139**, 151, 157
- Chemical indicators 72, 305, 307–309, 325
- Cirques 162, 173
- Code (Model) 330, 334, 336, 337
- Coefficient of efficiency 152, 240, 283, 342
- Conceptualisation of processes 15, 51–55, **53**, 63–70, **67**, **68**, 268–269, 330, 336, 366
 - see also* Model names
- Conceptual models 54
- Connectivity of patterns 10, 245–246
- Contour based terrain representation 56–59, **57**, 58, 80, 221–223, **222**, 258, **258**
- Convective storm 9, 127, 131
- Correlation length 12, 79
 - see also* Range
- Correlations
 - spatial – *see* Geostatistics
 - temporal 94, **96**, 103
- Coupling
 - land – atmosphere 84, 108–109, 122, 272
 - saturated – unsaturated zone 304–305, 310, 312

- Darcy's Law 65, 69, 222, 282
- Data – *see* Measurements
- Data assimilation 145–151, **148**, **150**, 157
- Deep seepage 220, 221, 237, 258–259, 313
- Degrees of freedom (model parameters) 72, 75, 153, 282, 337, 342–343
- Depth area duration methods (precipitation) 82, 100–101, **101**, 140

- Depth to water table 193, *193*, 221–222, 284, 309, *311*, 311–312, 316
- Dielectric constant 18, 135
- Digital elevation models 3–4, 14, 39–40
 accuracy 39–40, 56–59, 61, 277
 attribute computation 39–40, 57–59, 171, 224, 275–277
 grid size 39–40, 63
 scale effects 39–40, 61–62, 277
 types of 56–59, 57, 58
- Digital terrain models – *see* Digital elevation models
- Direct insertion (data assimilation) 145
- Disaggregation 28, 37–38
see also Downscaling
- Discharge – *see* Runoff
- Discharge zones – *see* Groundwater
- Discretisation of models 27, 56–74, 153, 282–283, 314
- Distributed parameter models 55, 63–65, 129, 164–167
see also Model names
- Distribution models
 rainfall-runoff 59, 63, 66, 70, 210, 272–273
 snow 59, 166
- Dominant processes 11, 14, 63, 248, 366
- Downscaling 84, 92–97, *93*, *94*, *95*, 341
see also Disaggregation
- Downward approach to modelling 71–72
- Drainage network 6, *6*, 153, 188, 199–200, 202, 320–321
- Dual-source approaches (evaporation) 114, 118–120
- Dunne mechanism – *see* Saturation excess runoff
- Dynamic scaling (precipitation) 88–92, 94, *94*, 98–100, *100*
- Eddy correlation, eddy covariance 109, 110–111
- Effective parameters 38, *67*, 68–69, *69*, 117, 166, 184
- Energy balance
 in evaporation 105, 106, 108, 119, 127
 in snowmelt 159–163, 171, 178–181
- Environmental management 4, 15, 125, 330, 353–354
- Ephemeral streams 119, 127, *128*, 138
- Equifinality of models – *see* Non-uniqueness
- Equilibrium evaporation 108
- Equilibrium modelling 304–305, 310–313, *311*, 324
- Erosion 72, 152, 352
- Errors 23, 32, 33, 41, *42*, 77, 184, 286
- ERS1/2 194, 196, *196*, 198
- ESTAR 136, 140
- Evaporation/evapotranspiration 105
 bare soil 118, 313
 estimation of 109–112, 117–120
 governing factors 11, 105–107, 112
 measurement of 109–112, 120–121, 127
 patterns – *see* Observed patterns;
 Simulated patterns
 potential 127, 223, 310–311, 314
- Exfiltration *53*, 221, 280, 282, 287, 314
- Extent (scale) 18–22, 27
- External drift kriging (geostatistics) 36, 43–45, *44*
- Feedback mechanisms 108–109, 122, 158, 162, 305, 317
- Field capacity 221–222, 281
- Filtering (scales) 19–20, *20*, 85, 197–198
see also Interpolation; Smoothing
- Flow depth 59, 68
- Flow detectors *251*, 253–254
- Flow path length 64–65, *65*
- Footprint (remote sensing) 19, 111–112
see also Support
- Four dimensional data assimilation – *see* Data assimilation
- GCMs (General Circulation Models) 15, 64, 102
- Geostatistics (methods) 31–39
- GIS (Geographic Information Systems) 3, 28, 45, 63
- GLUE methodology 297–298
- GOES 118
- Gravimetric method (soil moisture) 134, 217
- Grid-based terrain representation 56–59, *57*, 58, 170–171
- Ground-truthing (remote sensing) 24, 136–138, *140*, 195–197, 332, 341
- Groundwater
 flow 65, 274, 305, 321–322
 model 310–312
 observation wells – *see* Piezometers
 recharge/discharge 304, 307–309, 315, 321–322
see also Deep seepage
- Heterogeneity
 of land surfaces 112–117, 122, 129

- of soils 25, 152, 247–248, **255**, 260, 361–362
- representation of 12–14, 68–70, 260, 269–270, 284
- scales of 113–114, 116
- see also* Correlation length
- Homogeneity 12, 89, 114, 154, 323
- Hortonian runoff – *see* Infiltration excess runoff
- HRUs (Hydrological Response Units) 61
- Hydraulic conductivity
 - effective 38, 68–69, **69**
 - measurement of 22, 226, 250–253, 274
 - pedo-transfer functions 40–41, **144**, 244, 316, 332, 343
 - saturated 247, 253, **253**, **255**, 315–316, 315, 316
 - transmissivity 273, 280–281, 284, **291**
- Hydrochemical measurements 255–257, 272, 307–309, 325
- Hydrograph separation 256–257, **257**

- Incidence angle (remote sensing) 202, **202**
- Infiltration 70, 151, 187, 248, 310–311
- Infiltration excess runoff 10, 69, 151, 218, 248, 255
- Initial conditions of models 155, **155**, 209, 224, 248
- Interflow – *see* Subsurface stormflow
- Intermittent streams – *see* Ephemeral streams
- Interpolation 26–28
 - artifacts **29**, 30, **35**, 36–37, 43–45, **44**, 61, 277–278
 - change of scale 27, 31, 38, 228
 - choice of methods 28, **29**, 33–36, 41–50, **44**, 277–278, 365
 - filter 27, 31, 43
 - methods 28–39, 130
 - use of surrogate data 27–28, 35–38, 41–45, 166
- Inverse distance interpolation 29, **29**, 136

- Kinematic wave 64–65, 151, 221, 258, 282
- KINEROS 59, 70, 151–154
- Kriging (geostatistics) 33, 36, 41–45, **44**, 79, 130–131

- LAI (Leaf area index) 143, **144**
- Landsat 119, 203
- Landscape units 116, 129, 250, 260
- Land surface schemes 15, 69, 305
- Land use change 15, 72, 352

- Lapse rate (atmospheric) 163
- Latent heat 106, 109, 119–120, **120**, 159, 163
- Lateral redistribution 64–67, 116–117, 226, 305–306
- LOWTRAN7 136, 160, 161
- Lumped models 55, 342

- Macropores, macropore flow 7–9, **8**, **53**, 69, 242, 255–256, 268, 270, 275, 294
- Manning's 'n' 64–65, 68–69, 151, 259, 260
- Measurements
 - accuracy of 18, 23–26, 217, 332
 - design – *see* Sampling design
 - errors in 23–26, 30, 32, 33, 41, 48–49, 79, 111–112, 133, 154, 297, 330
 - scale 11, 18–22, **21**, 27, 46, 211–212, 228, 341, 347
- Micro-topography 14, 67, 70
- Microwave 24, 134–138, 188, 194–197
- MIKE SHE 345
- Model elements
 - representation of sub-grid variability – *see* Sub-grid variability
 - size of 61–63, 153, 166, 277
- Modelling philosophy 12, 51–55, **54**, 63, 70–74, **73**, 302–303, 322–323, 334–342, **335**, **338**, 345–346, 353–354
- Model names
 - KINEROS 59, 70, 151–154
 - MIKE SHE 345
 - MODFLOW 310–312
 - SHE 11
 - Stanford watershed model 54
 - TAPES-C 224
 - Thales 221–223, **222**
 - TOPLATS 141–144
 - TOPMODEL 59, 66, 280–283
 - TOPOG 249, 257–259
 - UEB 178–181
 - VUTS 170–171
- Models
 - calibration of 75–78, 152–153, 283–286, 288–291, 336, 338, 346
 - choice of 70–74
 - complexity of 55, 63–67, 70–74, **73**, 149, 153, 167, 282–283
 - credibility of 338, 341–345, 362–363
 - development of 52–53, **54**, 337–339, 345–346
 - performance of 73–74, **148**, 229–230, 241, 263, 270–271, 291, 294, 320
 - scales of 27, 62–63, 277, 330

- Models (*cont.*)
 spatial structure of 27, 56–74, 153, 282–283, 314
- Model testing 75–78, 152–153, 182, 263, 287–291, 301–302, 329–331, 334–342, 347–354, 363–366
- Model types 12, 51–55, 210–211, 273
- Model users 28, 36, 45, 49–50, 51, 330, 340, 349, 353–354
- MODFLOW 310–312
- Monsoon '90 119, 136
- Monte Carlo methods 77, 296
- MOS (Monin-Obukhov Similarity) theory 110, 113, 115, **116**
- Moving polynomials (interpolation) **29**, 30
- Multi-criteria testing 72, 77, 299, 348
- Multitemporal patterns 80–81, 188–189, 195–196, 202
- Neutron moisture meter 217
- NEXRAD 83
- Non-uniqueness (models) 71–72, 296, 331, 339
- Nudging (data assimilation) 145–146
- Nugget (geostatistics) 31–32, **32**, 79, 217, 230
- Number of sampling points 26, 48–49, 130, 177, 193, 216, 252, 253–254, 275, 307
- Numerical weather prediction models 88, 97
see also Precipitation, models of
- Objective functions 75
see also Performance criteria
- Observed patterns 357–359
 evaporation 120, **120**
 macropores 7, **8**
 piezometric surface 5, 278–280, **279**
 precipitation **86**, **93**, 95, **95**, **135**, **139**, 140–141, **142**
 recharge/discharge 305, 307, **308**
 runoff 138–140, **138**, **252**, 264–268, **267**
 saturated source areas 5, **60**, 187, 193, **195**, 205–206, **220**
 snow 4, 171–175, **172**, 178, **180**, **183**
 soil moisture 5, 7, 41–42, **42**, **137**, **144**, 218, **219**, **236**
- Organisation of variability 5, 7, 85–88, 202–203, 211–212
 degrees of **7**, **13**, 22, 245, 247–248, 252, 260, **266**, 267
 imposed 12–14, **13**, 45, 66, 75–76, 247–248, 294, 342–343
- Orographic precipitation 36, 163
- Overfitting **29**, 30, 339, 341
- Overland flow 67–68, **67**, 70, 151, 223, 254, 256, 264–269
- Parameterisation
 of precipitation 83–84, 87, 102
 of runoff processes 70, 143, 225–226, 258–259, 274
- Parameters
 effective 38, **67**, 68–69, **69**, 117, 166, 184
 identifiability 71–73, **73**, 286, 290–291, 331, 339
 optimisation of 75–78, 152–153, 283–286, 288–291, 336, 338, 346
 scale effects 72
- Particle size distribution 41, 274
- Pattern comparison methods
 optimal local alignment 80
 point by point 78–79, **120**, 173–174, **176**, 182, **185**, 229–230, **294**, 319–320, **320**
 residual patterns 78–79, 230–234, **233**
 stratification by terrain attributes 79, **176**, 246, 287, **288**
 visual 5, 78, 182, 228, 267, 289, 317–321
- Patterns
 binary 168, **172**, **175**, **195**, **220**, 361
 complementarity of patterns and time series 361–362
 observed – *see* Observed patterns
 relationship to process 5–11, **6**, 87, 136, 203, 234–238, 245, 356–361
 scale effects 11, 18–22, 114, 211–212, **212**
 simulated – *see* Simulated patterns
 summary of use of patterns 356–364, 357–359
 to identify model parameters 240–246, 267–268, 285–286, 315–321, 330
 to identify model structure 95–97, 149, 171–175, 182, 199–200, 227–238, 240–246, 285–296, 330
 utility of 3–5, 73, 97–98, 185–186, 244–246, 269, 271, 324–325, 329–331, 354, 356, 357–359
- PCA (Principal Component Analysis) 200–206, **202–204**, **206**
- Pedo-transfer functions 40–41, **144**, 244, 316, 332, 343
- Penman-Monteith model (evaporation) 107–108, 114, 118, 224
- Performance criteria (models) 74–81, 336, 337, 339, 341–342, 346
- Philip equation (infiltration) 311

- Physically based models 54, 151, 165–166, 178, 210–211, 221–223, 249, 257–259, 342
- Piezometer 217, 275, 305
- Pixel size 19, 114, 170, 194
see also Model elements
- Pollution 55, 72
- Porous media flow 52, 65, 69
- Precipitation
areal reduction of 82, 140
depth area duration methods 82, 100–101, **101**, 140
distribution, coverage 70, 82, 131–132, **132**, **133**
duration 82, 131–132
fluctuations 85, 89, 98
intensity 131–132, 248, 252
measurement of 21, 103, 127, 130–131
models of 84, 100–101, 140–141
patterns – *see* Observed patterns;
Simulated patterns
persistence of 96
scales of 9, **10**, 40, 85, 90–91, **91**, 100, **100**, 133, 163–164
types of 9
see also Convective storm; Orographic precipitation
variability of 82–83, 88–92, **96**, 129–134, **134**, 156–157
- Preferential flow 7–9, **8**, 52, **53**, 242, 256, 268
- Priestley-Taylor equation (evaporation) 108, 115, 118
- Proxy basin testing 340
- Proxy data – *see* Surrogate data
- Pushbroom radiometer 135, **137**, 143, **144**, 145, 156
- RADAR (precipitation) 24, 83, 85, **86**
see also SAR
- Radiation
diffuse 161
longwave 161–162, 173, 178–181
net 112, 121, 178
shortwave/global 119, 159–162, 178–181
- Radiosondes 102, 111
- Rainfall – *see* Precipitation
- Rainfall-runoff models – *see* Model names
- Rain gauges – *see* Precipitation, measurement of
- Range (geostatistics) 31–32, **32**
- Rate limiting processes 306
- REA (Representative Elementary Area) 62
- Realisations (geostatistics) 37, **44**, 260
- Recharge 304, 307–309, 315, 321–322
see also Deep seepage
- Regularisation (geostatistics) 20
- Remote sensing 16, 332, 365
aerial photographs 165, 169, **174**
calibration 24, 136–138, **140**, 195–197, 332, 341
evapotranspiration 108, 112, 118–121
microwave 24, 134–138, 188, 194–197
precipitation 24, 83, 85, **86**
see also ESTAR
snow 169–170
soil moisture 21, 134–138, **144**, 188, 213
vegetation 115, 200, 203–204, **204**
- Resolution 18–22, 27, 31, 34–35, **35**
see also Model elements, size of
- Rill flow 14, 67, 70
- Riparian zone 189, 205
- Root zone 106, 115, 149, 312
- Roughness
aerodynamic 110, 114
Manning's n 64–65, 68–69, 141, 259, 260
surface 69, 195
- Runoff
measurement of 127, **128**, 253–254, 274, **275**
models of – *see* Model names
patterns – *see* Observed patterns;
Simulated patterns
processes – *see* Infiltration excess runoff;
Saturation excess runoff; Subsurface stormflow; Groundwater; Overland flow; Channel flow; Variable source area
routing 64–65, 152, 223
variability of 138–140, **138**, 156, 253–254, 261
- Runon infiltration 10, 220, 221
- Sampling design 11, 17–18, 21–22, 24–25, 45–50, 168–170, 177, 211–213, 242
- SAR (Synthetic Aperture Radar) 24, 188, 194–197, 341–342
- Satellite – *see* Remote sensing
- Saturated area – *see* Observed patterns;
Simulated patterns
- Saturation deficit
of atmosphere 106
of soil 151, 213, 238, **239**, 281
- Saturation excess runoff 9, 40, 69, 209, 221, 248, 255

Scale

- effects on patterns 11, 18–22, 87, 114, 211, 212, **212**
 - invariance 83
 - of evaporation 116
 - of groundwater flow 315
 - of measurement 11, 18–22, **21**, 27, 46, 211–212, 228, 341, 347
 - of precipitation 9, **10**, 40, 85, 90–91, **91**, 100, **100**, 133, 163–164
 - of processes (general) 9, **10**, 46, 157, 164, 217, 310
 - of runoff **10**, 156
 - of soil moisture 136, 245
 - see also* Digital elevation models; Heterogeneity; Interpolation; Modelling; Parameters; Remote sensing
 - triplet 18–22, **19**, 27, 46, 62–63, 211
- Scaling exponent 87, 98, **99**
- Semi-arid catchments 111, 125–127, 157
- Sensible heat 108, 109, 159, 163
- Sensitivity analysis 77, 143, 154–156, **154**, **155**, 182–184, 260, 315–321, 339
- Sequential Gaussian simulation (geostatistics) 37
- Sequential indicator simulation (geostatistics) 37
- SHE 11
- Sheetflow 59, 67–68, 70
 - see also* Overland flow
- Sill (geostatistics) 31–32, **32**
- Similarity theory – *see* MOS
- Simulated patterns 357–359
 - evaporation 120, **120**
 - piezometric surface 292, **295**
 - precipitation **93**, 95, **95**, **142**
 - recharge/discharge 315–321, **316–318**, 322–323
 - runoff 264–268, **266**
 - saturated source areas **60**, **195**, 198–206, **203**, **206**
 - snow 171–175, **172**, **175**, **183**
 - soil moisture 41–45, 143–144, **144**, 148–149, 150, 227–238, **231**, **236**, 240–246
- Sky view factor (radiation) 160–161
- Sloughing (snow) 164
- Smith and Parlange equation (infiltration) 151
- Smoothing 19, **20**, 31, 41–43, **42**, 49, 79, 228, **232**

Snow

- accumulation 159, 163–164, 178
- course 24, 170
- density 177, 178, **180**
- drift 167, **179**, 182–184
- drift factor 167, 171, 173, 181, **182**
- melt 11, 159, 162, 165, 312
- models of 164–166, 170–171, 178–181
- patterns – *see* Observed patterns; Simulated patterns
- redistribution 11, 163–164
- stakes 168
- tubes 177
- vegetation effects 162, 163, 164, 167
- water equivalent 167, 177, **183**
- wind effects 11, 162–163, 164, 167, 173, 178, **179**

Soil

- bulk density 134, 225–226
- depth 193, 213–215, 227, 235–237, 250–252, 260, 277, **278**
- heat flux 106, 107, 159
- heterogeneity 25, 152, 247–248, **255**, 260, 361–362
- macropores 7–9, **8**, **53**, 69, 242, 255–256, 268, 270, 275
- particle size distribution 41, 274
- porosity 143, 225–226, 260, 281, 285, **289**
- structure 41, 215
- texture 127, 143, 190, 193, 215, 250–252, 306, 312, **315**, **316**
- water content – *see* Soil moisture

Soil moisture

- measurement (ground based) 21, 23, 47, 134–138
 - see also* Observed patterns
 - measurement (remote sensing) 21, 134–138, **144**, 188, 213
 - patterns – *see* Observed patterns; Simulated patterns
 - seasonality 7, 47, 49, **219**, 230
 - storage 217, 226, 258–259, 282, 310–311
 - temporal evolution 49, **148**, 189, 199–200, 217, 238, **239**
 - topographic effects 7, 49, 218, 230–238
- Spacing (scale) 18–22, 27, 31, 34–35, **35**
- Specific humidity 107
- Speckle filter (remote sensing) 197–198
- SPI (Saturation Potential Index) **195**, 199–200, **201**
- Splines **29**, 30–31, 33, 36, 130–131, 230
- Split sample testing 76, 152, 181–182, 340
- Stability function (atmosphere) 110

- Stanford watershed model 54
- Statistical correction (data assimilation) 145
- Statistical interpolation (SI) (data assimilation) 146–147
see also Kriging
- Stochastic models 54–55, 310–312, 324
- Stochastic simulation (geostatistics) 37
- Stomatal conductance 106–107, 114
- Storage deficit 151, 213, 238, **239**, 281
- Storm – *see* Precipitation
- Streamtube approach – *see* Contour-based terrain representation
- Structural analysis (geostatistics) 32–33
- Sub-grid variability 67–70, 83, 92–93, 152, 166
- Subsurface stormflow **53**, 65, 222, 255–256
- Super observations (data assimilation) 147
- Support (scale) 18–22, 27, 31, 228
- Surface runoff 12–14, 238–240
see also Overland flow
- Surface temperature 106, 114, 118, **137**, 162, 178–181
- Surrogate data 18, 23–25, 40, **42**, **44**, 46, 307, 310, 332, 343, 365
 pedo-transfer functions 40–41, **144**, 244, 316, 332, 343
- SVAT (Soil vegetation atmosphere transfer) models 118, 121
- TAPES-C 224
- TDAS (Terrain data acquisition system) 215, **216**
- TDR (Time Domain Reflectometry) 21, **21**, 23–24, 48, 134, 215–217
- Temperate catchments 9, 47, 187, 190, 213, 344
- Terrain analysis
 indices 25, 40, 43, 171, 223
see also Digital elevation models
see also Wetness index
- Testing of models – *see* Model testing
- Thales 221–223, **222**
- Thiessen method (interpolation) 29, **29**, 43, **44**, 343
- TIN (Triangulated Irregular Network) 39, 56–59, **57**, 58, 322
- TOPLATS 141–144
- TOPMODEL 59, 66, 280–283
- TOPOG 249, 257–259
- Topography
 effects on backscattering (remote sensing) 188, 200–203
 hydrological effects of 7–12, 25, 36, 40, 160–162, 173, 218, 274, 318
 relationship to soil properties 41, 203, 218, 253
 representation of – *see* Digital elevation models
 shading by 10–11, 143, 160, 171, 181
- Transmissivity
 atmospheric 160, 181
 hydraulic 273, 280–281, 284, **291**
- Trend 19, **20**, 30, 133
- Tropical catchments 249–254
- Turbulent transport 106, 109–111, 163
- Turning Bands Method (geostatistics) 37
- UEB 178–181
- Uncertainty 296–301
 bounds 256–257, 298–301, **299**, **300**
 GLUE 297–298
 representation of 77–78
- Unsaturated zone 305–306, **311**
- Upscaling **93**, 167, 341
see also Aggregation
- Upward approach to modelling 71–72
- Vadose zone 305–306, **311**
- Validation – *see* Model testing
- Vapour pressure 107
- Variable source area 187, 188–189, 272
- Variogram (geostatistics) 31–32, **32**, 34, 136
- Vegetation
 effect on snow 162–163, 164, 167
 heterogeneity 112–117, 122, 129
 remote sensing of 115, 200, 203–204, **204**
- Verification – *see* Model testing
- Vertical redistribution 64–65, 217, 305–306
- VUTS 170–171
- Water content – *see* Soil moisture
- Water table 215, 272, 274; *see also* Depth to water table
 patterns – *see* Observer patterns;
 Simulated patterns
- Wavelet transform 85
- Weighting function (data assimilation) 146–147
- Wetness index 12–13, 187, 200
 computation of 39–40, 57–59, 224, 275–277
 quasi-dynamic 66
 steady state 40, 43, 59, 70, 144, **144**, 272–273, 280
- Wind
 drift factors 167, 171, 173, 181, **182**
 effect on evaporation 106, 115

Wind (*cont.*)

- effect on precipitation 133
- effect on snow patterns 11, 162–163, 164, 167, 173, 178, **179**

Zero plane displacement (atmospheric boundary layer) 115